

OPTIMAL MODEL SELECTION FOR DENSITY ESTIMATION OF STATIONARY DATA UNDER VARIOUS MIXING CONDITIONS.

BY MATTHIEU LERASLE*

* *Institut Mathématiques de Toulouse (UMR 5219), INSA Toulouse.*

We propose a block-resampling penalization method for marginal density estimation with non necessary independent observations. When the data are β or τ -mixing, the selected estimator satisfies oracle inequalities with leading constant asymptotically equal to 1. We also prove in this setting the slope heuristic, which is a data-driven method to optimize the leading constant in the penalty.

1. Introduction. Model selection by penalization of an empirical loss is a general method that includes several famous procedures as cross-validation (Rudemo (1982)) or hard thresholding (Donoho *et al.* (1996)) as shown by Barron, Birgé and Massart (1999). The difficulty is to calibrate the penalty so that the selected estimator satisfies an oracle inequality. A good penalty has the shape of an ideal one, see the definition (2.4), and depends in general on a leading constant that should be chosen sufficiently large.

Resampling penalties provide a shape for the penalty term in a general statistical learning framework, see Arlot (2009). The resulting estimator satisfies sharp oracle inequalities in non-Gaussian heteroscedastic regression among histograms (Arlot (2009)) and in density estimation among more general collections of models (Lerasle (2009b)). The validity of these theorems relies on the independence of the observation. In this paper, we study a generalization of these penalties, called block-resampling penalties and we prove that the resulting estimator satisfies sharp oracle inequalities when the data are only supposed to be β - or τ -mixing (the coefficient β has been defined by Volkonskiĭ and Rozanov (1959), the coefficient τ by Dedecker and Prieur (2005), see Section 2.4).

We use a coupling method to extend the results for independent data. It was introduced in Baraud, Comte and Viennet (2001) in a regression problem and used in Comte and Merlevède (2002) for density estimation with β -mixing observations. β is a well known "strong" mixing coefficient. We

AMS 2000 subject classifications: Primary 62G07, 62G09; secondary 62M99

Keywords and phrases: Density estimation, optimal model selection, resampling methods, slope heuristic, weak dependence

refer to the books of [Doukhan \(1994\)](#) and [Bradley \(2007\)](#) for examples of β -mixing processes. One of the most important is the following: a stationary, irreducible, aperiodic and positively recurrent Markov chain is β -mixing. "Strong" mixing coefficients cannot be used to study a lot of simple processes. For example [Andrews \(1984\)](#), the stationary solution of the equation

$$(1.1) \quad X_n = \frac{1}{2}(X_{n-1} + \xi_n),$$

where $(\xi_n)_{n \in \mathbb{Z}}$ are i.i.d. Bernoulli random variables $\mathcal{B}(1/2)$ is not β -mixing. This is why "weak" mixing coefficients as τ , have been introduced. They are easier to compute and allow to cover more examples, as the process (1.1) (see [Dedecker and Prieur \(2005\)](#), [Comte, Dedecker and Taupin \(2008\)](#) or the book [Dedecker et al. \(2007\)](#) for examples of weakly-mixing processes). In [Lerasle \(2009a\)](#), we used a coupling result of [Dedecker and Prieur \(2005\)](#) to extend the coupling method to τ -mixing data.

In these papers, the dimension of the models was used as a shape of the penalties, but the leading constant was built with the mixing coefficients. They cannot in general be computed from the data and even if they are, the theoretical upper bounds obtained are probably too pessimistic to be used by the statistician. The slope algorithm allows to calibrate this leading constant in an optimal way. It is based on the slope heuristic, introduced in [Birgé and Massart \(2007\)](#) and proved in [Birgé and Massart \(2007\)](#) for Gaussian regression, in [Arlot and Massart \(2009\)](#) for non-Gaussian heteroscedastic regression over histograms and in [Lerasle \(2009b\)](#) for density estimation. The second main result of this paper is a proof of the slope heuristic for the marginal density estimation problem with β - or τ -mixing data.

Block-resampling penalties and the slope heuristic can be defined in a more general statistical learning framework, including the problems of classification and regression (this framework is the one of [Massart and Nédélec \(2006\)](#)). Our results are contributions to the theoretical understanding of these generic methods. Up to our knowledge, they are the first ones obtained in a mixing framework.

The paper is organized as follows. Section 2 introduces the density estimation framework, the estimators, the penalties and the main assumptions. Sections 3 and 4 give the main results, respectively for τ - and β -mixing processes. Section 5 gives the proofs of the main results. Some other proofs are available as supplementary material ([Supplement A](#)).

2. Preliminaries.

2.1. The density estimation framework. We observe n real valued, identically distributed random variables $X_1, \dots, X_n$, defined on a probability space

$(\Omega, \mathcal{A}, \mathbb{P})$, with common law P . We assume that P is absolutely continuous with respect to the Lebesgue measure μ on $\mathbb{R}$ and we want to estimate the density s of P with respect to μ . $L^2(\mu)$ denotes the Hilbert space of square integrable real valued functions and $\|\cdot\|$ the associated L^2 -norm. We assume that s belongs to $L^2(\mu)$. The risk of an estimator $\hat{s}$ of s is measured with the L^2 -loss, that is $\|s - \hat{s}\|^2$, which is random when $\hat{s}$ is.

Let p, q be two integers and assume that $n = 2pq$. For all $i = 0, \dots, p-1$, let $I_i = (2iq + 1, \dots, (2i+1)q)$, $A_i = (X_l)_{l \in I_i}$. For all functions t in $L^1(P)$, for all reals $x_1, \dots, x_q$, we define

$$L_q t(x_1, \dots, x_q) = \frac{1}{q} \sum_{i=1}^q t(x_i), \quad P t = \int_{\mathbb{R}} t(x) s(x) d\mu(x), \quad P_A t = \frac{1}{p} \sum_{i=0}^{p-1} L_q t(A_i).$$

Given a linear space S_m of measurable, real valued functions, and an orthonormal basis $(\psi_\lambda)_{\lambda \in \Lambda_m}$ of S_m , we define the projection estimator $\hat{s}_{A,m}$ of s onto S_m by

$$\hat{s}_{A,m} = \sum_{\lambda \in \Lambda_m} (P_A \psi_\lambda) \psi_\lambda = \arg \min_{t \in S_m} \{ \|t\|^2 - 2P_A t \}.$$

Given a finite collection $(S_m)_{m \in \mathcal{M}_n}$ of such linear spaces and a penalty function $\text{pen} : \mathcal{M}_n \rightarrow \mathbb{R}_+$, the Penalized Projection Estimator, hereafter PPE is defined by

$$(2.1) \quad \tilde{s}_A = \hat{s}_{A, \hat{m}}, \text{ where } \hat{m} \in \arg \min_{m \in \mathcal{M}_n} \{ \|\hat{s}_{A,m}\|^2 - 2P_A \hat{s}_{A,m} + \text{pen}(m) \}.$$

We will say that the PPE satisfies an oracle inequality when one of the two following inequalities holds.

There exist constants $\kappa > 0, \gamma > 1$ and a positive sequence $(K_n)_{n \in \mathbb{N}^*}$ bounded away from 0 such that

$$(2.2) \quad \mathbb{P} \left(K_n \|s - \tilde{s}_A\|^2 \leq \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) \geq 1 - \frac{\kappa}{n^\gamma}.$$

There exists a positive sequence $(K_n)_{n \in \mathbb{N}^*}$ bounded away from 0 such that

$$(2.3) \quad K_n \mathbb{E} (\|s - \tilde{s}_A\|^2) \leq \mathbb{E} \left(\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right).$$

The oracle inequality is said to be sharp when, moreover, the sequence $K_n \rightarrow 1$ when n grows to infinity. Inequalities (2.2) are usually preferred to (2.3) since they describe the typical behavior of the selected estimator and not only of its expectation.

It is worth mentioning that we only use $\text{Card}(\cup_{i=0}^{p-1} I_i) = pq = n/2$ data to build the estimator $\tilde{s}_A$. The consequences of this choice are discussed after Theorem 3.1 and in Section 4.3.

2.2. Block-resampling penalties. We introduce block-resampling penalties as natural generalizations of resampling penalties. The best estimator in the collection $(\hat{s}_{A,m})_{m \in \mathcal{M}_n}$ minimizes among $\mathcal{M}_n$ the ideal criterion

$$\|s - \hat{s}_{A,m}\|^2 - \|s\|^2 = \|\hat{s}_{A,m}\|^2 - 2P_A \hat{s}_{A,m} + \text{pen}_{\text{id}}(m).$$

In this decomposition, the ideal penalty $\text{pen}_{\text{id}}(m)$ (Arlot (2009)) is equal to

$$(2.4) \quad \text{pen}_{\text{id}}(m) = 2(P_A - P)(\hat{s}_{A,m}).$$

To adapt the approach of Arlot (2009) to a dependent setting, we replace the resampling step by a resampling procedure on the blocks $(A_i)_{i=0,\dots,p-1}$. Let $(W_0, \dots, W_{p-1})$ be a resampling scheme, that is, a vector of positive random variables, independent of $(X_i)_{i=1,\dots,n}$ and exchangeable, which means that, for all permutations ξ of $\{0, \dots, p-1\}$,

$$(W_{\xi(0)}, \dots, W_{\xi(p-1)}) \text{ has the same law as } (W_0, \dots, W_{p-1}).$$

Let $\overline{W} = p^{-1} \sum_{i=0}^{p-1} W_i$, for all t in $L^1(P)$, let P_A^W be the block-resampling empirical process defined by

$$P_A^W t = \frac{1}{p} \sum_{i=0}^{p-1} W_i L_q t(A_i).$$

For all integrable random variables $F(X_1, \dots, X_n, W_0, \dots, W_{p-1})$, let

$$\mathbb{E}_W[F(X_1, \dots, X_n, W_0, \dots, W_{p-1})] = \mathbb{E}[F(X_1, \dots, X_n, W_0, \dots, W_{p-1}) | X_1, \dots, X_n].$$

Let $((\psi_\lambda)_{\lambda \in \Lambda_m})_{m \in \mathcal{M}_n}$ be orthonormal bases of $(S_m)_{m \in \mathcal{M}_n}$ and let $(\hat{s}_{A,m}^W)_{m \in \mathcal{M}_n}$ be the collection of resampling projection estimators

$$\hat{s}_{A,m}^W = \sum_{\lambda \in \Lambda_m} (P_A^W \psi_\lambda) \psi_\lambda.$$

The block-resampling penalties are defined as block-resampling estimators of the ideal penalty by

$$(2.5) \quad \text{pen}_W(m, C) = C \mathbb{E}_W (2(P_A^W - \overline{W} P_A)(\hat{s}_{A,m}^W)).$$

The idea of resampling is to mimic the behavior of the empirical process P_A around P by the behavior of the resampling empirical process P_A^W around $\overline{W} P_A$. The resampling procedure is a plug-in method where the unknown functionals $F(P, P_n)$ are estimated by $F(\overline{W} P_n, P_n^W)$. Hence, $\hat{s}_{A,m}$

in $\text{pen}_{\text{id}}(m)$ is replaced by $\widehat{s}_{A,m}^W$ in $\text{pen}_W(m, C)$ and, instead of applying the process $P_A - P$, we apply the process $P_A^W - \overline{W}P_A$. We take the expectation with respect to the distribution of the resampling scheme to stabilize the procedure. Finally, we let a normalizing constant C free for this general definition.

We use a block-resampling scheme instead of a classical exchangeable resampling scheme in order to preserve the dependence of the data inside the blocks. This is a key point for the procedure to work. Examples of resampling schemes can be found in [Arlot \(2009\)](#). The classical block-bootstrap, [Künsch \(1989\)](#); [Liu and Singh \(1992\)](#) is obtained when the distribution of $(W_0, \dots, W_{p-1})$ is the multinomial $\mathcal{M}(p, 1/p, \dots, 1/p)$.

2.3. The Slope Algorithm. The "slope heuristic" has been introduced by [Birgé and Massart \(2007\)](#) in order to calibrate the leading constant in a penalty term (for example the constant C in (2.5)). It is based on the behavior of the complexity of the selected model (recall the definition (2.1)). It states that there exist a family $(\Delta_m)_{m \in \mathcal{M}_n}$ and a constant $K_{\min}$ satisfying the following properties.

- SH1 When $\text{pen}(m) \leq K\Delta_m$, with $K < K_{\min}$, then $\Delta_{\hat{m}} \geq c_1 \max_{m \in \mathcal{M}_n} \Delta_m$.
- SH2 When $\text{pen}(m) = K\Delta_m$, with $K > K_{\min}$, then $\Delta_{\hat{m}}$ is much smaller.
- SH3 When $\text{pen}(m) = 2K_{\min}\Delta_m$, then $\widehat{s}_A$ satisfies an optimal oracle inequality.

Based on this heuristic, [Birgé and Massart \(2007\)](#) introduced the following slope algorithm. It can be used in practice when a family $(\Delta_m)_{m \in \mathcal{M}_n}$ satisfying the slope heuristic is known.

- For all $K > 0$, compute $\Delta_{\hat{m}(K)}$ where $\hat{m}(K)$ is defined as in (2.1) with $\text{pen}(m) = K\Delta_m$.
- Find $\tilde{K}$ such that $\Delta_{\hat{m}(K)}$ is very large for $K < \tilde{K}$ and much smaller when $K > \tilde{K}$.
- Choose the final $\hat{m}$ equal to $\hat{m}(2\tilde{K})$.

The idea is that $\tilde{K} \sim K_{\min}$ since we observe a jump of the complexity of the selected model around $K = \tilde{K}$ (thanks to **SH1**, **SH2**) and thus that the final estimator, selected by the penalty $2\tilde{K}\Delta_m \sim 2K_{\min}\Delta_m$, satisfies an optimal oracle inequality (by **SH3**).

2.4. Some measures of dependence.

2.4.1. β -mixing data. [Volkonskiĭ and Rozanov \(1959\)](#) defined the coefficient β as follows. Let Y be a random variable defined on a probability space

$(\Omega, \mathcal{A}, \mathbb{P})$ and let $\mathcal{M}$ be a σ -algebra in $\mathcal{A}$, let

$$\beta(\mathcal{M}, \sigma(Y)) = \mathbb{E} \left(\sup_{A \in \mathcal{B}} |\mathbb{P}_{Y|\mathcal{M}}(A) - \mathbb{P}_Y(A)| \right).$$

For all stationary sequences of random variables $(X_n)_{n \in \mathbb{Z}}$ defined on $(\Omega, \mathcal{A}, \mathbb{P})$, let

$$\beta_k = \beta(\sigma(X_i, i \leq 0), \sigma(X_i, i \geq k)).$$

The process $(X_n)_{n \in \mathbb{Z}}$ is said to be β -mixing when $\beta_k \rightarrow 0$ as $k \rightarrow \infty$.

2.4.2. τ -mixing data. [Dedecker and Prieur \(2005\)](#) defined the coefficient τ as follows. For all l in $\mathbb{N}^*$, for all x, y in $\mathbb{R}^l$, let $d_l(x, y) = \sum_{i=1}^l |x_i - y_i|$. For all l in $\mathbb{N}^*$, for all functions t defined on $\mathbb{R}^l$, the Lipschitz semi-norm of t is defined by

$$\text{Lip}_l(t) = \sup_{x \neq y \in \mathbb{R}^l} \frac{|t(x) - t(y)|}{d_l(x, y)}.$$

For all functions t defined on $\mathbb{R}$, we will denote for short by $\text{Lip}(t) = \text{Lip}_1(t)$. Let λ_1 be the set of all functions $t : \mathbb{R}^l \rightarrow \mathbb{R}$ such that $\text{Lip}_l(t) \leq 1$. For all integrable, $\mathbb{R}^l$ -valued, random variables Y defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ and all σ -algebra $\mathcal{M}$ in $\mathcal{A}$, let

$$\tau(\mathcal{M}, Y) = \mathbb{E} \left(\sup_{t \in \lambda_1} |\mathbb{P}_{Y|\mathcal{M}}(t) - \mathbb{P}_Y(t)| \right).$$

For all stationary sequences of integrable random variables $(X_n)_{n \in \mathbb{Z}}$ defined on $(\Omega, \mathcal{A}, \mathbb{P})$, for all integers k, r , let

$$\tau_{k,r} = \max_{1 \leq l \leq r} \frac{1}{l} \sup_{k \leq i_1 < \dots < i_l} \{ \tau(\sigma(X_p, p \leq 0), (X_{i_1}, \dots, X_{i_l})) \}, \quad \tau_k = \sup_{r \in \mathbb{N}^*} \tau_{k,r}.$$

The process $(X_n)_{n \in \mathbb{Z}}$ is said to be τ -mixing when $\tau_k \rightarrow 0$ as $k \rightarrow \infty$.

2.5. Main Assumptions.

2.5.1. A specific collection for τ -mixing sequences. Wavelet spaces have been widely used in density estimation, in particular, because the oracle is adaptive over Besov spaces (see [Birgé and Massart \(1997\)](#)).

Dyadic Wavelet spaces: Let r be a real number, $r \geq 1$. We work with an r -regular orthonormal multiresolution analysis of $L^2(\mu)$, associated with a compactly supported scaling function ϕ and a compactly supported mother wavelet ψ . Without loss of generality, we suppose that the support of the

functions ϕ and ψ is included in an interval $[A_1, A_2)$ where A_1 and A_2 are integers such that $A_2 - A_1 = A \geq 1$. For all k in $\mathbb{Z}$ and j in $\mathbb{N}^*$, let $\psi_{0,k} : x \rightarrow \sqrt{2}\phi(2x - k)$ and $\psi_{j,k} : x \rightarrow 2^{j/2}\psi(2^j x - k)$. The family $\{(\psi_{j,k})_{j \geq 0, k \in \mathbb{Z}}\}$ is an orthonormal basis of $L^2(\mu)$. The collection of dyadic wavelet spaces is described as follows.

[W] *dyadic wavelet generated spaces:* let $J_n = [\log_2(n)]$, for all $J_m = 1, \dots, J_n$, let

$$\Lambda_m = \{(j, k), 0 \leq j \leq J_m, k \in \mathbb{Z}\}$$

and let S_m be the linear span of $\{\psi_\lambda\}_{\lambda \in \Lambda_m}$.

2.5.2. General framework: We present in this section a set of hypothesis sufficient to prove the theorems. None of them is used to build the penalties.

H1 : *There exists a constant κ_a such that, for all m, m' in $\mathcal{M}_n$, for all t in $S_m + S_{m'}$, with $\|t\| \leq 1$, there exist t_m in S_m and $t_{m'}$ in $S_{m'}$, with $\|t_m\| \vee \|t_{m'}\| \leq \kappa_a$ such that $t = t_m + t_{m'}$.*

This assumption is typically satisfied for nested collections as **[W]**.

H2 : $N_n = \text{Card}(\mathcal{M}_n)$ is finite and there exist constants $c_{\mathcal{M}}, \alpha_{\mathcal{M}}$ such that $N_n \leq c_{\mathcal{M}} n^{\alpha_{\mathcal{M}}}$.

This assumption means that the collection is not too rich and thus, that the model selection problem is not too hard. It is satisfied by the collection **[W]**.

Let us introduce some notations. For all m in $\mathcal{M}_n$, for all orthonormal bases $(\psi_\lambda)_{\lambda \in \Lambda_m}$ of S_m , let

$$D_{A,m} = q \sum_{\lambda \in \Lambda_m} \text{Var}(L_q(\psi_\lambda)(A_0)), \quad R_{A,m} = n \|s - s_m\|^2 + 2D_{A,m},$$

$$B_m = \{t \in S_m, \|t\| \leq 1\}, \quad b_m = \sup_{t \in B_m} \|t\|_\infty.$$

$D_{A,m}$, and thus $R_{A,m}$, are well defined since we can check with Cauchy-Schwarz inequality that

$$D_{A,m} = q \mathbb{E} \left[\left(\sup_{t \in B_m} L_q t(A_0) - P t \right)^2 \right].$$

Two quantities will play a fundamental role to discuss the results. The first one is the risk of an oracle.

$$R_n = \inf_{m \in \mathcal{M}_n} R_{A,m}.$$

We are typically interested in non parametric problems where $R_n/n \sim n^{-\gamma}$ for some $0 < \gamma < 1$. This situation occurs, for example, when s is a regular function, in this case, we have $R_n/n = \kappa n^{-2\alpha/(2\alpha+1)}$, for some $\alpha > 0$, $\kappa > 0$. We will make the following assumption.

H3 : *There exists a constant $\kappa_R > 0$ such that $R_n \geq \kappa_R (\ln n)^8$.*

The constant 8 in $(\ln n)^8$ is technical, it yields the rate $\varepsilon_n = (\ln n)^{-1/2}$ in the oracle inequalities. [Arlot \(2009\)](#) replaced this assumption by a lower bound on the bias of the models. It implies that $R_n \geq \kappa n^\gamma$, for some constants $\kappa > 0$, $1 > \gamma > 0$ and therefore Assumption **H3**.

H4 : *There exists a constant $c_D > 0$ such that,*

$$\forall m \in \mathcal{M}_n, P \left(\sup_{t \in B_m} t^2 \right) \geq c_D b_m^2.$$

It is shown in Section 6 that some classical examples of collections $(S_m, m \in \mathcal{M}_n)$ as regular histograms, Fourier spaces and **[W]** satisfy **H4**.

The following assumptions will be used to prove the slope heuristic. We introduce a second quantity, that will play a fundamental role. Let

$$D_n^* = \max_{m \in \mathcal{M}_n} D_{A,m}.$$

In classical collection of models, like **[W]**, $D_n^* \sim cn$. This is why we introduce the following assumption.

H5 : *$D_n^*/R_n \rightarrow \infty$ when n grows to infinity.*

We will prove that, when the data are mixing $D_{A,m} \simeq n \mathbb{E} (\|s_m - \widehat{s}_{A,m}\|^2)$ represents the variance term of the risk. It is a natural measure of the complexity of the models. Hence, D_n^* represents the maximal complexity of the models. Moreover, R_n is the risk of the oracle. It balances the complexity and the bias term and has therefore the same order as the complexity of an oracle. Hence, Assumption **H5** means that the largest complexity in the collection $(S_m)_{m \in \mathcal{M}_n}$ is much larger than the one of an oracle, which is a natural condition for the slope heuristic to hold. We need a final assumption.

H6 : *For all m^* such that $D_{A,m^*} = D_n^*$, we have*

$$\frac{n \|s - s_{m^*}\|^2}{D_n^*} \rightarrow 0 \text{ when } n \rightarrow \infty.$$

When D_n^* is of order n , **H6** simply means that the distance between s and a complex model goes to 0. In general, it means that for these complex

models, the bias part of the risk is negligible compared to the variance part. We conclude this section by the assumptions on the mixing coefficients. All mean that these coefficients are sufficiently small. Let $\gamma = \beta$ or τ .

[AR(θ)] *arithmetical γ -mixing with rate θ : there exists $C > 0$ such that, for all k in $\mathbb{N}$, $\gamma_k \leq C(1+k)^{-(1+\theta)}$.*

S(β) : $\sum_{l \geq 1} (l+1)\beta_l \leq c_D/64$, where c_D is defined in **H4**.

We prove in Appendix that $c_D = 1$ for regular histograms and Fourier spaces.

S($\tau, \mathbf{W}$) : $\sum_{l \geq 1} \left(\|s\|^2 \tau_l \right)^{1/3} \leq C(W)$, where $C(W)$ depends only on ϕ, ψ .

The value of the constant $C(W)$ is given in the supplementary material [Supplement A](#).

3. Results for τ -mixing sequences.

3.1. Resampling penalties. The result of this section is that PPE selected by block-resampling penalties satisfy sharp oracle inequalities.

THEOREM 3.1. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of regular wavelet spaces **[W]** satisfying **H3, H4**. Let p, q be two integers such that $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$.*

Let $\tilde{C}_W = \text{Var}(W_1 - \bar{W})^{-1}$, $C > \tilde{C}_W/2$ and let $\tilde{s}_A$ be the PPE defined in (2.1) with the penalty $\text{pen}_W(m, C)$ defined in (2.5).

*Assume that there exists $\theta > 5$ such that $X_1, \dots, X_n$ are arithmetically **[AR(θ)]** τ -mixing and satisfy **S($\tau, \mathbf{W}$)**. Let $\varepsilon_n = (\ln n)^{-1/2}$, $\kappa(C) = 2 \left(C\tilde{C}_W^{-1} - 1 \right)$.*

There exist constants κ_1, κ_2 such that we have

$$(3.1) \quad K_n \mathbb{E}(\|s - \tilde{s}_A\|^2) \leq \mathbb{E} \left(\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) + \frac{\kappa_2}{n},$$

with

$$K_n = \frac{(1 \wedge (1 + \kappa(C))) - \kappa_1 \varepsilon_n}{(1 \vee (1 + \kappa(C))) + \kappa_1 \varepsilon_n}.$$

Comments:

- The constant C has to be chosen asymptotically equal to $\tilde{C}_W$. However, from a non-asymptotic point of view, it is helpful to choose C slightly larger. A first reason is that (3.1) is useless when $K_n \leq 0$, which occurs if $\kappa_1 \varepsilon_n > 1$ and $C = \tilde{C}_W$.

3.2. Slope heuristic. Theorem 3.1 gives a totally data driven penalty which satisfies a sharp oracle inequality, therefore, the heuristic is not necessary to obtain asymptotically optimal results. However, we saw that C can be optimized for small samples. Moreover, the slope algorithm is faster to compute than the resampling penalties when a deterministic quantity can be used in the slope heuristic. Theorem 3.2 hereafter justifies property **SH1** of the heuristic. Δ_m is the variance term $D_{A,m}/n$ and $K_{\min} = 2$.

THEOREM 3.2. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of regular wavelet spaces **[W]** satisfying **H3**, **H4**, **H5**, **H6**. Let p, q be two integers such that $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$. Assume that there exists a constant $0 < \delta < 1$ such that , for all m in $\mathcal{M}_n$,*

$$(3.2) \quad 0 \leq \text{pen}(m) \leq (2 - \delta) \frac{D_{A,m}}{n},$$

and let $\tilde{s}_A$ be the PPE defined in (2.1).

Assume that there exists $\theta > 5$ such that $X_1, \dots, X_n$ are arithmetically **[AR](θ)** τ -mixing and satisfy **S**($\tau, \mathbf{W}$). There exist constants κ_1, κ_2 such that

$$(3.3) \quad \mathbb{E}(D_{A,\hat{m}}) \geq \frac{4\delta}{9} D_n^* - \kappa_1.$$

$$(3.4) \quad \mathbb{E}(\|s - \tilde{s}_A\|^2) \geq \frac{\delta}{5} \frac{D_n^*}{R_n} \left(\mathbb{E} \left(\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) - \frac{\kappa_2}{n} \right).$$

Comments:

- Inequality (3.3) states that $D_{A,\hat{m}}$ is as large as possible when the penalty term is too small. This is exactly **SH1** with $\Delta_m = D_{A,m}$.
- Inequality (3.4) states that the model selected by a too small penalty is never an oracle. This is another reason why it is interesting to choose $C > \tilde{C}_W$ in Theorem 3.1.

The following theorem justifies properties **SH2**, **SH3** of the slope heuristic.

THEOREM 3.3. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of regular wavelet spaces **[W]** satisfying **H3**, **H4**. Let p, q be two integers such that $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$. Assume that there exist $\delta_+ \geq -\delta_- > -1$, $\varepsilon \geq 0$ and some constants κ_1, κ_2*

satisfying, for all m in $\mathcal{M}_n$,

$$(3.5) \quad \mathbb{E} \left[\sup_{m \in \mathcal{M}_n} \left((2 - \delta_-) \frac{2D_{A,m}}{n} - \text{pen}(m) - \varepsilon \frac{R_{A,m}}{n} \right)_+ \right] \leq \frac{\kappa_1}{n},$$

$$(3.6) \quad \mathbb{E} \left[\sup_{m \in \mathcal{M}_n} \left(\text{pen}(m) - (2 + \delta_+) \frac{D_{A,m}}{n} - \varepsilon \frac{R_{A,m}}{n} \right)_+ \right] \leq \frac{\kappa_2}{n}.$$

Let $\tilde{s}_A$ be the PPE defined in (2.1) with $\text{pen}(m)$ and let $\varepsilon_n = (\ln n)^{-1/2}$.

Assume that there exists $\theta > 5$ such that $X_1, \dots, X_n$ are arithmetically $[\mathbf{AR}(\theta)]$ τ -mixing and satisfy $\mathbf{S}(\tau, \mathbf{W})$. There exist constants $\kappa_1, \kappa_2, \kappa_3$, such that

$$(3.7) \quad K_n \mathbb{E} (\|\tilde{s}_A - s\|^2) \leq \mathbb{E} \left(\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) + \frac{\kappa_2}{n},$$

with

$$K_n = \frac{(1 \wedge (1 - \delta_-)) - \kappa_1(\varepsilon_n + \varepsilon)}{(1 \vee (1 + \delta_+)) + \kappa_1(\varepsilon_n + \varepsilon)}.$$

Moreover, we have

$$(3.8) \quad K_n \mathbb{E} (D_{A,\hat{m}}) \leq R_n + \kappa_3.$$

Comments:

- When $\text{pen}(m)$ becomes larger than $2D_{A,m}/n$, $D_{A,\hat{m}}$ jumps from D_n^* (3.3) to R_n ((3.8) for δ_+ and $-\delta_-$ close to -1). This justifies **SH2** since $R_n < D_n^*$.
- A model selected with a penalty $4D_{A,m}/n$ satisfies an oracle inequality (Theorem 3.3 for δ_+ and δ_- close to 0). This justifies **SH3**.
- $D_{A,m}$ is unknown and cannot be used in the slope algorithm. It can be shown (see Lemma 5.2 in the supplementary material) that $D_{A,m}$ satisfies $\kappa_* 2^{J_m} \leq D_{A,m} \leq \kappa^* 2^{J_m}$. The slope heuristic might hold for $\Delta_m = 2^{J_m}/n$, but a complete proof requires moreover that $\kappa_* \simeq \kappa^*$. However, we obtain in the proof of Theorem 3.1 that, for $C = \tilde{C}_W$, $\text{pen}_W(m, C)$ satisfies (3.5) and (3.6) for $\delta_+ = \delta_- = 0$ and $\varepsilon = \kappa \varepsilon_n$. Since (3.2) can be modified to work with random penalties, we can apply the slope algorithm with $\text{pen}_W(m, 1)$ instead of $D_{A,m}/n$.

4. Results for β -mixing sequences. We show that block-resampling penalties select oracles and that the slope heuristic holds in this case.

4.1. Resampling penalties.

THEOREM 4.1. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of linear spaces satisfying **H1**, **H2**, **H3**, **H4**. Let p, q be two integers such that $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$.*

Let $\tilde{C}_W = \text{Var}(W_1 - \bar{W})^{-1}$, $C > \tilde{C}_W/2$ and let $\tilde{s}_A$ be the PPE defined in (2.1) with the block-resampling penalty $\text{pen}_W(m, C)$ defined in (2.5).

Assume that there exists $\theta > 2$ such that $X_1, \dots, X_n$ are arithmetically $[\mathbf{AR}(\theta)]$ β -mixing and satisfy $\mathbf{S}(\beta)$. Let $\varepsilon_n = (\ln n)^{-1/2}$, $\kappa(C) = 2 \left(C\tilde{C}_W^{-1} - 1 \right)$.

There exist constants κ_1, κ_2 such that,

(4.1)

$$P \left(K_n \|s - \tilde{s}_A\|^2 \leq \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) \geq 1 - \kappa_2 \left(\frac{1}{n^2} \vee \frac{(\ln n)^{4+2\theta}}{n^{\theta/2}} \right),$$

with

$$K_n = \frac{(1 \wedge (1 + \kappa(C))) - \kappa_1 \varepsilon_n}{(1 \vee (1 + \kappa(C))) + \kappa_1 \varepsilon_n}$$

Comments:

- The coupling lemma of [Berbee \(1979\)](#) for β -mixing processes is much stronger than the one satisfied by τ -mixing data ([Dedecker and Prieur \(2005\)](#)). This is why Theorem 4.1 covers more collections of models than Theorem 3.1 and why we prove oracle inequalities in probability.

4.2. *Slope heuristic.* The following theorems are adaptations to the β -mixing case of Theorems 3.2 and 3.3.

THEOREM 4.2. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of linear spaces satisfying **H1**, **H2**, **H3**, **H4**, **H5**, **H6**. Let p, q be two integers such that $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$.*

Let $\tilde{s}_A$ be the PPE defined in (2.1) with a penalty $\text{pen}(m)$ satisfying, for all m in $\mathcal{M}_n$, Condition (3.2) of Theorem 3.2.

Assume that there exists $\theta > 2$ such that $X_1, \dots, X_n$ are arithmetically $[\mathbf{AR}(\theta)]$ β -mixing and satisfy $\mathbf{S}(\beta)$. There exists a constant κ and an event Ω_n such that

$$\mathbb{P}(\Omega_n) \geq 1 - \kappa \left(\frac{1}{n^2} \vee \frac{(\ln n)^{4+2\theta}}{n^{\theta/2}} \right),$$

and, on Ω_n ,

$$(4.2) \quad D_{A,\hat{m}} \geq \frac{4\delta}{9} D_n^*, \quad \|s - \tilde{s}_A\|^2 \geq \frac{\delta}{5} \frac{D_n^*}{R_n} \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2.$$

THEOREM 4.3. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of linear spaces satisfying **H1**, **H2**, **H3**, **H4**. Let p, q be two integers such that $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$. Assume that there exist $\delta_+ \geq -\delta_- > -1$, $\varepsilon \geq 0$, $0 \leq \eta < 1$ and an event Ω_{pen} , with $\mathbb{P}(\Omega_{pen}) \geq 1 - \eta$ such that, on Ω_{pen} , for all m in $\mathcal{M}_n$,*

$$(4.3) \quad (2 - \delta_-) \frac{2D_{A,m}}{n} - \varepsilon \frac{R_{A,m}}{n} \leq pen(m) \leq (2 + \delta_+) \frac{2D_{A,m}}{n} + \varepsilon \frac{R_{A,m}}{n}.$$

Let $\tilde{s}_A$ be the PPE defined in (2.1) with pen .

Assume that there exists $\theta > 2$ such that $X_1, \dots, X_n$ are arithmetically $[\mathbf{AR}(\theta)]$ β -mixing and satisfy $\mathbf{S}(\beta)$. There exist constants κ_1, κ_2 and an event Ω_n^* such that

$$\mathbb{P}(\Omega_n^*) \geq 1 - \eta - \kappa_2 \left(\frac{1}{n^2} \vee \frac{(\ln n)^{4+2\theta}}{n^{\theta/2}} \right),$$

and, on Ω_n^* ,

$$(4.4) \quad K_n \|\tilde{s}_A - s\|^2 \leq \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2,$$

with

$$K_n = \frac{(1 \wedge (1 - \delta_-)) - \kappa_1(\varepsilon_n + \varepsilon)}{(1 \vee (1 + \delta_+)) + \kappa_1(\varepsilon_n + \varepsilon)}.$$

Moreover, Ω_n^* , $2K_n D_{A,\hat{m}} \leq 3R_n$.

Comments:

- We refer to the comments of Theorems 3.2 and 3.3 where we explain why Theorems 4.2 and 4.3 imply the slope heuristic with $\Delta_m = D_{A,m}/n$, $K_{\min} = 2$.
- As in Theorem 3.3, $D_{A,m}$ cannot be used to build a model selection procedure. A deterministic shape of $D_{A,m}$ is unknown, although we prove in the supplementary material that $D_{A,m}$ is bounded by b_m^* . However, $pen_W(m, 1)$ can be used instead of $D_{A,m}$.

4.3. Discussion and perspectives. Block-resampling penalties are data driven procedures for the estimation of the marginal density of mixing data. They satisfy optimal oracle inequalities without remainder term. The theorems hold for possibly infinite dimensional models. Finally the oracle inequalities are sharp and there is no remainder term. This improves Theorems 3.1 and 4.1 in Lerasle (2009a) and Theorem 3.1 in Comte and Merlevède (2002).

Lacour (2008) gave a model selection procedure to estimate the stationary density and the transition probability of a Markov Chain. She worked with a stationary chain, irreducible, aperiodic and positively recurrent, which is therefore β -mixing. Her density estimator is selected by a penalty equal to Kd_m/n with a constant K that "depends on the law of the chain" (see Remark 4 after Theorem 3 in Lacour (2008)) and that she proposed to estimate in the simulations by the slope algorithm. We prove the slope heuristic, justifying that the slope algorithm can be used to optimize the leading constant. It would be interesting to see if resampling penalties may be used in her context to estimate the transition probabilities.

Gannaz and Wintenberger (2009) worked with other weak mixing coefficients (namely λ and ϕ , see Dedecker *et al.* (2007) for a definition) and studied a wavelet thresholded estimator. The main advantage is that the thresholded estimator is adaptive over a larger class of Besov spaces than the oracle over the collection $[\mathbf{W}]$ (for details about this important issue see Barron, Birgé and Massart (1999)). The main drawback is that their threshold is built with the mixing coefficients.

Block-resampling penalties can be extended to the statistical learning framework of Massart and Nédélec (2006), where the slope algorithm has already been defined (Arlot and Massart (2009)). We believe that these procedures perform well in this context but the problem remains open.

The main drawback of our approach is that we use only $n/2$ data. Moreover, the deterministic choice of the number p of blocks is not optimized. For example, when the data are geometrically β -mixing, which means that, for some constants $\theta > 0$, $C > 0$, $\beta_k \leq Ce^{-\theta k}$, choosing p of order $n(\ln n)^{-2}$ would improve the rates of convergence of the leading constant. An interesting direction of research would be to provide data-driven choices of p and q to improve these rates, and a data-driven choice of blocks to use more data. In practice, the computation time is also a very important issue. Actually, the conditional expectation is a bit long to evaluate and some efforts have to be done in this direction. Things can be improved if we obtain a deterministic shape of the ideal penalty, as in the independent case, since the slope heuristic is faster to compute with a deterministic Δ_m . We obtain upper and lower bounds on pen_{id} , but our inequalities are not sharp enough to justify completely the slope heuristic. We can also think of the V -fold cross validation penalties defined in Arlot (2008). These penalties are also faster to compute than the resampling penalties. They can be viewed as resampling penalties defined with non-exchangeable weights. These issues are far beyond the objectives of the present paper and will be addressed in forthcoming works.

Acknowledgements: The author is very grateful to Béatrice Laurent and Clémentine Prieur, for their precious advices. He would like also to thank the reviewers and associate editors for their careful reading of the manuscript and helpful comments which led to an improved presentation of the paper.

5. Proofs.

5.1. *Notations.* Recall that p and q are integers such that $2pq = n$, and that $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$. For all $k = 0, \dots, p-1$, let $I_k = (2kq + 1, \dots, (2k+1)q)$, $A_k = (X_i)_{i \in I_k}$ and $I = \cup_{k=0}^{p-1} I_k$. For all t in $L^2(\mu)$ and all $x_1, \dots, x_q$ in $\mathbb{R}$,

$$L_q(t)(x_1, \dots, x_q) = \frac{1}{q} \sum_{i=1}^q t(x_i), \quad P_A t = \frac{1}{p} \sum_{k=0}^{p-1} L_q(t)(A_k) = \frac{2}{n} \sum_{i \in I} t(X_i),$$

$$\nu_A(t) = (P_A - P)(t).$$

For all m in $\mathcal{M}_n$, we denote by $(\psi_\lambda)_{\lambda \in \Lambda_m}$ an orthonormal basis of S_m . The estimator $\widehat{s}_{A,m}$ associated to the model S_m , is defined as

$$\widehat{s}_{A,m} = \sum_{\lambda \in \Lambda_m} (P_A \psi_\lambda) \psi_\lambda.$$

Classical computations show that, if s_m denotes the orthogonal projection of s onto S_m ,

$$s_m = \sum_{\lambda \in \Lambda_m} (P \psi_\lambda) \psi_\lambda, \text{ hence } \|\widehat{s}_{A,m} - s_m\|^2 = \sum_{\lambda \in \Lambda_m} (\nu_A \psi_\lambda)^2.$$

The ideal penalty, $2\nu_A(\widehat{s}_{A,m})$ satisfies

$$\nu_A(\widehat{s}_{A,m} - s_m) + \nu_A(s_m) = \sum_{\lambda \in \Lambda_m} (\nu_A \psi_\lambda)^2 + \nu_A(s_m) = \|\widehat{s}_{A,m} - s_m\|^2 + \nu_A(s_m).$$

For all m, m' in $\mathcal{M}_n$, let

$$p(m) = \|s_m - \widehat{s}_{A,m}\|^2 = \sup_{t \in B_m} (\nu_A(t))^2 = \sum_{\lambda \in \Lambda_m} (\nu_A(\psi_\lambda))^2.$$

$$\delta(m, m') = 2\nu_A(s_m - s_{m'}).$$

Hereafter $W_0, \dots, W_{p-1}$ denotes a resampling scheme, $\overline{W} = p^{-1} \sum_{i=0}^{p-1} W_i$, P_A^W denotes the resampling empirical process, defined for all measurable functions t by

$$P_A^W t = \frac{1}{p} \sum_{i=0}^{p-1} W_i L_q t(A_i).$$

We introduce also $\nu_A^W = P_A^W - \overline{W} P_A$ and $\tilde{C}_W = (\text{Var}(W_1 - \overline{W}))^{-1}$. For any orthonormal basis $(\psi_\lambda)_{\lambda \in \Lambda_m}$ of S_m , let

$$p_W(m) = \tilde{C}_W \sum_{\lambda \in \Lambda_m} \mathbb{E}_W ((\nu_A^W(\psi_\lambda))^2).$$

$p_W(m)$ is well defined since, from Cauchy-Schwarz inequality,

$$p_W(m) = \tilde{C}_W \mathbb{E}_W \left(\sup_{t \in B_m} (\nu_A^W t)^2 \right).$$

Let $\varepsilon_n = (\ln n)^{-1/2}$ and let $\kappa > 0$. Let $\mathcal{M}$ denote one of the set $\mathcal{M}_n$ or $\mathcal{M}_n^2$. When $\mathcal{M} = \mathcal{M}_n$, for all $\overline{m}$ in $\mathcal{M}$ let $R_{A,\overline{m}} = R_{A,m}$ and when $\mathcal{M} = \mathcal{M}_n^2$, for all $\overline{m} = (m, m')$ in $\mathcal{M}$, let $R_{A,\overline{m}} = R_{A,m} \vee R_{A,m'}$. For all $\overline{m}$ in $\mathcal{M}$, let

$$(5.1) \quad f_1(\overline{m}, \kappa) = p(m) - \frac{2D_{A,m}}{n} - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(5.2) \quad f_2(\overline{m}, \kappa) = \frac{2D_{A,m}}{n} - p(m) - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(5.3) \quad f_3(\overline{m}, \kappa) = p(m) - p_W(m) - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(5.4) \quad f_4(\overline{m}, \kappa) = p_W(m) - p(m) - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(5.5) \quad f_5(\overline{m}, \kappa) = \delta(m, m') - \kappa \varepsilon_n \frac{R_{A,m} \vee R_{A,m'}}{n}.$$

We will use the following fact.

Fact 0: The resampling penalty $\text{pen}_W(m, C)$ defined in (2.5) satisfies

$$\text{pen}_W(m, C) = 2C\tilde{C}_W^{-1}p_W(m).$$

Proof: Let $(\psi_\lambda)_{\lambda \in \Lambda_m}$ be an orthonormal basis of S_m . Recall that $\hat{s}_{A,m}^W = \sum_{\lambda \in \Lambda_m} (P_A^W \psi_\lambda) \psi_\lambda$, so that

$$\hat{s}_{A,m}^W - \overline{W} \hat{s}_{A,m} = \sum_{\lambda \in \Lambda_m} (\nu_A^W \psi_\lambda) \psi_\lambda.$$

Hence, $\nu_A^W(\hat{s}_{A,m}^W - \overline{W}\hat{s}_{A,m}) = \sum_{\lambda \in \Lambda_m} (\nu_A^W \psi_\lambda)^2$.

We conclude the proof showing that $\mathbb{E}_W(\nu_A^W(\overline{W}\hat{s}_{A,m})) = 0$, hence

$$\frac{p_W(m)}{\tilde{C}_W} = \mathbb{E}_W(\nu_A^W(\hat{s}_{A,m}^W - \overline{W}\hat{s}_{A,m})) = \mathbb{E}_W(\nu_A^W(\hat{s}_{A,m}^W)) = \frac{\text{pen}_W(m, C)}{2C}.$$

Since $W_0, \dots, W_{p-1}$ are independent of $X_1, \dots, X_n$,

$$\mathbb{E}_W(\nu_A^W(\overline{W}\hat{s}_{A,m})) = \frac{1}{p^2} \sum_{i,j=0}^{p-1} L_q(\psi_\lambda)(A_i) L_q(\psi_\lambda)(A_j) \mathbb{E}_W(W_i \overline{W} - (\overline{W})^2).$$

Then, by exchangeability of the weights,

$$\begin{aligned} \mathbb{E}_W(W_i \overline{W} - (\overline{W})^2) &= \frac{1}{p} \left(\mathbb{E}(W_i^2) + \sum_{j \neq i} \mathbb{E}(W_i W_j) \right) \\ &\quad - \frac{1}{p^2} \left(\sum_i \mathbb{E}(W_i^2) + \sum_{i \neq j} \mathbb{E}(W_i W_j) \right) = 0. \square \end{aligned}$$

5.2. *Proof of Theorem 3.1.* The proof is based on the following Lemma, whose proof is given in the additional material.

LEMMA 5.1. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of regular wavelet spaces $[\mathbf{W}]$ satisfying assumptions **H3, H4**. Let p, q be two integers satisfying $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$. Assume that there exists $\theta > 5$ such that $X_1, \dots, X_n$ are arithmetically $[\mathbf{AR}(\theta)]$ τ -mixing and satisfy $\mathbf{S}(\tau, \mathbf{W})$. There exist constants κ_1, κ_2 , such that, for all $i = 1, \dots, 5$, for all $\overline{m}$ in $\mathcal{M}$,*

$$(5.6) \quad \mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} (f_i(\overline{m}, \kappa_1))_+ \right) \leq \frac{\kappa_2}{n}.$$

It comes from Fact 0 and the equality $2C\tilde{C}_W = \kappa(C) + 2$ that, for all m in $\mathcal{M}_n$,

$$(5.7) \quad \text{pen}_W(m, C) - (2 + \kappa(C))p(m) = 2C\tilde{C}_W^{-1}(p_W(m) - p(m)).$$

Hence, from (5.6) with $i = 3, 4$, $\text{pen}_W(m, C)$ satisfies Conditions (3.6) and (3.5) of Theorem 3.3 with $\delta_+ = -\delta_- = \kappa(C)$ and $\varepsilon = 2\kappa_1 C \tilde{C}_W^{-1} \varepsilon_n$. Theorem 3.1 follows from (3.7).

5.3. *Proof of Theorem 4.1.* The proof is based on the following lemma whose proof is given in additional material.

LEMMA 5.2. *Let $\theta > 1$ and let $(X_n)_{n \in \mathbb{Z}}$ be an arithmetically $[\mathbf{AR}(\theta)]$ β -mixing process satisfying $\mathbf{S}(\beta)$. Let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of linear spaces satisfying Assumptions **H1**, **H2**, **H3**, **H4**. Let p, q such that $2pq = n$, $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$. There exist constants κ_1, κ_2 which may vary from line to line such that, on an event Ω_n satisfying*

$$\mathbb{P}(\Omega_n) \geq 1 - \kappa_2 \left(\frac{(\ln n)^{2(1+\theta)}}{n^{\theta/2}} \vee \frac{1}{n^2} \right), \text{ where}$$

$$(5.8) \quad \forall \bar{m} \in \mathcal{M}, \forall i = 1, \dots, 5, f_i(\bar{m}) \leq 0.$$

Hence, from (5.6) with $i = 3, 4$, $\text{pen}_W(m, C)$ satisfies Condition (4.3) of Theorem 4.3 with $\delta_+ = -\delta_- = \kappa(C)$ and $\varepsilon = 2\kappa_1 C \tilde{C}_W^{-1} \varepsilon_n$. Theorem 4.1 follows from (4.4).

5.4. *Proof of Theorem 3.2 and 4.2.* It is sufficient to prove the results for sufficiently large n since we can increase the constant κ_2 if necessary. Let m_o be a model such that $R_{A, m_o} = R_n$. Now, by definition, $\hat{m}$ minimizes among $\mathcal{M}_n$ the following criterion:

$$\text{Crit}(m) = \|\hat{s}_{A, m}\|^2 - 2P_A \hat{s}_{A, m} + \text{pen}(m) + \|s\|^2 + 2\nu_A(s_{m_o}).$$

Fact 1: For all m in $\mathcal{M}_n$,

$$\text{Crit}(m) = \|s_m - s\|^2 + \text{pen}(m) - p(m) + 2\nu_A(s_{m_o} - s_m).$$

Proof: Recalling that $\|s - \hat{s}_{A, m}\|^2 = \|\hat{s}_{A, m}\|^2 - 2P \hat{s}_{A, m} + \|s\|^2$ and that $(P_A - P)(\hat{s}_{A, m} - s_m) = \|\hat{s}_{A, m} - s_m\|^2 = p(m)$, we have,

$$\begin{aligned} \text{Crit}(m) &= \|s - \hat{s}_{A, m}\|^2 - 2\nu_A(\hat{s}_{A, m} - s_m) + 2\nu_A(s_{m_o} - s_m) + \text{pen}(m) \\ &= (\|s - \hat{s}_{A, m}\|^2 - \|\hat{s}_{A, m} - s_m\|^2) - p(m) + \text{pen}(m) + 2\nu_A(s_{m_o} - s_m). \end{aligned}$$

We conclude the proof with Pythagoras equality. $\square$

Fact 2: For all m in $\mathcal{M}_n$, for all constants κ_1 ,

$$\begin{aligned} (1 + 2\kappa_1 \varepsilon_n) \frac{2D_{A, m}}{n} &\geq -\text{Crit}(m) + (1 - 2\kappa_1 \varepsilon_n) \|s - s_m\|^2 \\ &\quad - \sup_{m \in \mathcal{M}_n} (f_1(m, \kappa_1)) - \sup_{(m, m') \in \mathcal{M}_n^2} (f_5((m, m'), \kappa_1)). \end{aligned}$$

Proof: From Fact 1, for all m in $\mathcal{M}_n$, for all κ_1 , since $\text{pen}(m) \geq 0$,

$$\text{Crit}(m) \geq \|s_m - s\|^2 - f_1(m, \kappa_1) - \frac{2D_{A,m}}{n} - 2\kappa_1\varepsilon_n \frac{R_{A,m}}{n} - f_5((m_o, m), \kappa_1).$$

We conclude the proof using that $R_{A,m} = n\|s - s_m\|^2 + 2D_{A,m}$. $\square$

Fact 3: For all m in $\mathcal{M}_n$, for all constants κ_1 ,

$$\begin{aligned} (\delta - 4\kappa_1\varepsilon_n) \frac{D_{A,m}}{n} &\leq -\text{Crit}(m) + (1 + 2\kappa_1\varepsilon_n) \|s - s_m\|^2 \\ &\quad + \sup_{m \in \mathcal{M}_n} (f_2(m, \kappa_1)) + \sup_{(m, m') \in \mathcal{M}_n^2} (f_5((m, m'), \kappa_1)). \end{aligned}$$

Proof: From Fact 1, for all m in $\mathcal{M}_n$, for all κ_1 , since $\text{pen}(m) \leq (2 - \delta)D_{A,m}/n$,

$$\text{Crit}(m) \leq \|s_m - s\|^2 + f_2(m, \kappa_1) - \delta \frac{D_{A,m}}{n} + 2\kappa_1\varepsilon_n \frac{R_{A,m}}{n} + f_5((m, m_o), \kappa_1).$$

We conclude the proof using that $R_{A,m} = n\|s - s_m\|^2 + 2D_{A,m}$. $\square$

From Fact 2, we have, for all κ_1 ,

$$\begin{aligned} (1 + 2\kappa_1\varepsilon_n) \frac{2D_{A,\hat{m}}}{n} &\geq -\text{Crit}(\hat{m}) + (1 - 2\kappa_1\varepsilon_n) \|s - s_{\hat{m}}\|^2 \\ &\quad - \sup_{m \in \mathcal{M}_n} (f_1(m, \kappa_1)) - \sup_{(m, m') \in \mathcal{M}_n^2} (f_5((m, m'), \kappa_1)). \end{aligned}$$

Let us now consider a model m^* such that $D_{A,m^*} = D_n^*$. By definition of $\hat{m}$, we have $\text{Crit}(\hat{m}) \leq \text{Crit}(m^*)$. Hence, from Fact 3, we deduce that,

$$\begin{aligned} (1 + 2\kappa_1\varepsilon_n) \frac{2D_{A,\hat{m}}}{n} &\geq -\text{Crit}(m^*) + (1 - 2\kappa_1\varepsilon_n) \|s - s_{\hat{m}}\|^2 \\ &\quad - \sup_{m \in \mathcal{M}_n} (f_1(m, \kappa_1)) - \sup_{(m, m') \in \mathcal{M}_n^2} (f_5((m, m'), \kappa_1)). \\ (5.9) \quad &\geq (\delta - 4\kappa_1\varepsilon_n) \frac{D_{A,m^*}}{n} + (1 - 2\kappa_1\varepsilon_n) \|s - s_{\hat{m}}\|^2 - (1 + 2\kappa_1\varepsilon_n) \|s - s_{m^*}\|^2 \\ &\quad - \sup_{(m, m') \in \mathcal{M}_n^2} (f_1(m, \kappa_1) + f_2(m, \kappa_1) + 2f_5((m, m'), \kappa_1)). \end{aligned}$$

From Lemma 5.1, there exist κ_1 and κ_2 such that

$$\mathbb{E} \left(\sup_{(m, m') \in \mathcal{M}_n^2} (f_1(m, \kappa_1) + f_2(m, \kappa_1) + 2f_5((m, m'), \kappa_1)) \right) \leq \frac{\kappa_2}{n}.$$

From Lemma 5.2, there exists κ_1 such that, on Ω_n ,

$$\sup_{(m,m') \in \mathcal{M}_n^2} (f_1(m, \kappa_1) + f_2(m, \kappa_1) + 2f_5((m, m'), \kappa_1)) \leq 0.$$

Now, assume that n is sufficiently large to ensure that

$$4\kappa_1\varepsilon_n \leq \frac{\delta}{4} \leq \frac{1}{4}, \quad \frac{n\|s - s_{m^*}\|^2}{D_n^*} \leq \frac{2\delta}{9}.$$

Then, taking the expectation in (5.9), we obtain that

$$\frac{9\mathbb{E}(D_{A,\hat{m}})}{8n} \geq \frac{\delta}{2} \frac{D_n^*}{n} - \frac{\kappa_4}{n}.$$

Hence (3.3) is proved for n sufficiently large.

Moreover, on Ω_n , we have

$$\frac{9D_{A,\hat{m}}}{8n} \geq \frac{\delta}{2} \frac{D_n^*}{n}.$$

Hence the first inequality of (4.2) is proved for n sufficiently large. (3.4) and the second inequality of (4.2) follow from the inequality

$$\|s - \tilde{s}_A\|^2 \geq (1 - \kappa_1\varepsilon_n) \frac{R_{A,\hat{m}}}{n} - f_2(\hat{m}, \kappa_1).$$

From Lemma 5.1, there exist constants κ_1, κ_2 , such that $\mathbb{E}(f_2(\hat{m}, \kappa_1)) \leq \kappa_2/n$. We choose n sufficiently large to ensure that $\kappa_1\varepsilon_n \leq 1/2$, we use (3.3) and we obtain that there exists a constant κ such that

$$\mathbb{E}(\|s - \tilde{s}_A\|^2) \geq \frac{2\delta}{9} \frac{D_n^* - \kappa}{n}.$$

We conclude the proof of (3.4) with the following Fact.

Fact 4:

$$\begin{aligned} \frac{R_n}{n} &\geq \frac{16}{17} \mathbb{E} \left(\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) - \frac{\kappa}{n}, \\ \text{thus } \frac{D_n^*}{n} &\geq \frac{16D_n^*}{17R_n} \left(\mathbb{E} \left(\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \right) - \frac{\kappa}{n} \right). \end{aligned}$$

Proof of Fact 4. Let κ_1 , be the constant previously defined,

$$\inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \leq (1 + \kappa_1\varepsilon_n) \inf_{m \in \mathcal{M}_n} \left\{ \frac{R_{A,m}}{n} \right\} + \sup_{m \in \mathcal{M}_n} f_1(m, \kappa_1).$$

We conclude the proof with Lemma 5.1. $\square$

We use the first inequality of (4.2) and we obtain that, on Ω_n ,

$$\|s - \tilde{s}_A\|^2 \geq \frac{2\delta}{9} \frac{D_n^*}{n}.$$

We conclude the proof of Theorem 4.2, saying that, on Ω_n , we have

$$\begin{aligned} \frac{R_n}{n} &= \inf_{m \in \mathcal{M}_n} \left\{ \|s - s_m\|^2 + \frac{2D_{A,m}}{n} \right\} \geq (1 - \kappa_1 \varepsilon_n) \inf_{m \in \mathcal{M}_n} \{ \|s - s_m\|^2 + p(m) \} \\ &= (1 - \kappa_1 \varepsilon_n) \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \geq \frac{15}{16} \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2. \end{aligned}$$

Thus,

$$\|\tilde{s}_A - s\|^2 \geq \frac{2\delta}{9} \frac{D_n^*}{R_n} \frac{R_n}{n} \geq \frac{\delta}{9} \frac{D_n^*}{R_n} \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2.$$

5.5. Proof of Theorems 3.3 and 4.3. As in the previous proof, it is sufficient to obtain the results for sufficiently large n . Let us first prove the oracle inequalities. Let κ_1 be a constant to be chosen later. Let Ω_n be the set defined on Lemma 5.2. The key point to prove oracle inequalities is the following fact.

Fact 5: For all m in $\mathcal{M}_n$, for all real numbers δ_- , δ_+ and for all non negative reals x, y ,

$$\begin{aligned} &[(1 \wedge (1 - \delta_-)) - x - y] \|s - \tilde{s}_A\|^2 \leq [(1 \vee (1 + \delta_+)) + x + y] \|s - \hat{s}_{A,m}\|^2 \\ (5.10) \quad &+ \sup_{m \in \mathcal{M}_n} \{ \text{pen}(m) - (2 + \delta_+) \|\hat{s}_{A,m} - s_m\|^2 - x \|s - \hat{s}_{A,m}\|^2 \}_+ \end{aligned}$$

$$(5.11) \quad + \sup_{m \in \mathcal{M}_n} \{ (2 - \delta_-) \|\hat{s}_{A,m} - s_m\|^2 - \text{pen}(m) - x \|s - \hat{s}_{A,m}\|^2 \}_+$$

$$(5.12) \quad + 2 \sup_{(m, m') \in \mathcal{M}_n^2} \{ \nu_A(s_{m'} - s_m) - y (\|s - \hat{s}_{A,m}\|^2 + \|s - \hat{s}_{A,m'}\|^2) \}_+.$$

Proof: By definition of $\tilde{s}_A$, for all m in $\mathcal{M}_n$, we have,

$$\|\tilde{s}_A\|^2 - 2P_A \tilde{s}_A + \text{pen}(\hat{m}) + \|s\|^2 \leq \|\hat{s}_{A,m}\|^2 - 2P_A \hat{s}_{A,m} + \text{pen}(m) + \|s\|^2.$$

Now, for all m in $\mathcal{M}_n$, since $\|\hat{s}_{A,m} - s\|^2 = \|\hat{s}_{A,m}\|^2 - 2P \hat{s}_{A,m} + \|s\|^2$,

$$\|\hat{s}_{A,m}\|^2 - 2P_A \hat{s}_{A,m} + \|s\|^2 = \|\hat{s}_{A,m} - s\|^2 - 2(P_A - P) \hat{s}_{A,m}.$$

Thus, for all m in $\mathcal{M}_n$,

$$\|\tilde{s}_A - s\|^2 - 2(P_A - P) \tilde{s}_A + \text{pen}(\hat{m}) \leq \|\hat{s}_{A,m} - s\|^2 - 2(P_A - P) \hat{s}_{A,m} + \text{pen}(m).$$

For all m in $\mathcal{M}_n$, since $(P_A - P)(\widehat{s}_{A,m} - s_m) = \|\widehat{s}_{A,m} - s\|^2$,

$$2(P_A - P)\widehat{s}_{A,m} = 2\|s_m - \widehat{s}_{A,m}\|^2 + 2(P_A - P)s_m.$$

This yields

$$\begin{aligned} \|s - \tilde{s}_A\|^2 &\leq \|s - \widehat{s}_{A,m}\|^2 + \text{pen}(m) - 2\|\widehat{s}_{A,m} - s_m\|^2 \\ &\quad + 2\|\widehat{s}_{A,\widehat{m}} - s_{\widehat{m}}\|^2 - \text{pen}(\widehat{m}) + 2\nu_A(s_{\widehat{m}} - s_m). \end{aligned}$$

We add $-(\delta_- \vee 0) + (x + y)\|\tilde{s}_A - s\|^2$ to the left hand side of the previous inequality and $-\delta_- \|\tilde{s}_A - s_{\widehat{m}}\|^2 - (x + y)\|\tilde{s}_A - s\|^2 + [(\delta_+ \vee 0) + x + y]\|s - \widehat{s}_{A,m}\|^2 - \delta_+ \|\widehat{s}_{A,m} - s_m\|^2 - (x + y)\|s - \widehat{s}_{A,m}\|^2$ to the right hand side. This is valid because, for all m in $\mathcal{M}_n$, for all reals δ ,

$$[(\delta \vee 0) + x + y]\|\widehat{s}_{A,m} - s\|^2 \geq \delta\|\widehat{s}_{A,m} - s_m\|^2 + (x + y)\|\widehat{s}_{A,m} - s\|^2.$$

We obtain

$$\begin{aligned} [(1 \wedge (1 - \delta_-)) - x - y]\|s - \tilde{s}_A\|^2 &\leq [(1 \vee (1 + \delta_+)) + x + y]\|s - \widehat{s}_{A,m}\|^2 \\ &\quad + \text{pen}(m) - (2 + \delta_+)\|\widehat{s}_{A,m} - s_m\|^2 - x\|\widehat{s}_{A,m} - s\|^2 \\ &\quad + (2 - \delta_-)\|\widehat{s}_{A,\widehat{m}} - s_{\widehat{m}}\|^2 - \text{pen}(\widehat{m}) - x\|\widehat{s}_{A,\widehat{m}} - s\|^2 \\ &\quad + 2\nu_A(s_{\widehat{m}} - s_m) - y\|\widehat{s}_{A,m} - s\|^2 - x\|\widehat{s}_{A,\widehat{m}} - s\|^2. \square \end{aligned}$$

We will also use the following fact.

Fact 6 : For all reals κ such that $\kappa\varepsilon_n \leq 1/2$,

$$\frac{R_{A,m}}{n} \leq 2\|s - \widehat{s}_{A,m}\|^2 + 2\{f_2(m, \kappa)\}_+.$$

Proof of Fact 6: We write

$$\frac{R_{A,m}}{n} = \frac{1}{1 - \kappa\varepsilon_n} \left(\frac{R_{A,m}}{n} - \|s - \widehat{s}_{A,m}\|^2 - \kappa\varepsilon_n \frac{R_{A,m}}{n} \right) + \frac{1}{1 - \kappa\varepsilon_n} \|s - \widehat{s}_{A,m}\|^2.$$

We use that $\kappa\varepsilon_n \leq 1/2$ and that $R_{A,m} = 2D_{A,m} + n\|s - s_m\|^2$ to conclude the proof. $\square$

Control of (5.10). Assume that n is sufficiently large to ensure that $\kappa_1\varepsilon_n \leq 1/2$. We have, from Fact 6,

$$\begin{aligned} \text{pen}(m) - (2 + \delta_+)p(m) - 2\varepsilon\|\widehat{s}_{A,m} - s\|^2 \\ \leq \text{pen}(m) - (2 + \delta_+)p(m) - \varepsilon \frac{R_{A,m}}{n} + 2\varepsilon\{f_2(m, \kappa_1)\}_+. \end{aligned}$$

Applying Lemma 5.1, we obtain constants κ_1 and κ_2 such that

$$\mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \{f_2(m, \kappa_1)\}_+ \right) \leq \frac{\kappa_2}{n}.$$

Applying Lemma 5.2, we obtain a constant κ_1 such that, on Ω_n ,

$$\sup_{m \in \mathcal{M}_n} \{f_2(m, \kappa_1)\}_+ \leq 0.$$

Moreover, (3.6) ensures that

$$\mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \left\{ \text{pen}(m) - (2 + \delta_+)p(m) - \varepsilon \frac{R_{A,m}}{n} \right\}_+ \right) \leq \frac{\kappa}{n}.$$

On Ω_{pen} we have

$$\sup_{m \in \mathcal{M}_n} \left\{ \text{pen}(m) - (2 + \delta_+)p(m) - \varepsilon \frac{R_{A,m}}{n} \right\}_+ \leq 0.$$

We choose $x = 2\varepsilon$. We obtain that, for Theorem 3.1, the expectation of (5.10) is upper bounded by κn^{-1} and for Theorem 4.1, the term (5.10) is equal to 0 on $\Omega_n \cap \Omega_{\text{pen}}$.

Control of (5.11). Assume that n is sufficiently large to ensure that $\kappa_1 \varepsilon_n < 1/2$, we deduce from Fact 6 that

$$\begin{aligned} & (2 - \delta_-)p(m) - \text{pen}(m) - 2\varepsilon \|\widehat{s}_{A,m} - s\|^2 \\ & \leq (2 - \delta_-)p(m) - \text{pen}(m) - \varepsilon \frac{R_{A,m}}{n} + 2\varepsilon \{f_2(m, \kappa_1)\}_+. \end{aligned}$$

Applying Lemma 5.1, we obtain constants κ_1 and κ_2 such that

$$\mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \{f_2(m, \kappa_1)\}_+ \right) \leq \frac{\kappa_2}{n}.$$

Applying Lemma 5.2, we obtain a constant κ_1 such that, on Ω_n ,

$$\sup_{m \in \mathcal{M}_n} \{f_2(m, \kappa_1)\}_+ \leq 0.$$

Moreover, (3.5) ensures that

$$\mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \left\{ (2 - \delta_-)p(m) - \text{pen}(m) - \varepsilon \frac{R_{A,m}}{n} \right\}_+ \right) \leq \frac{\kappa}{n}.$$

On Ω_{pen} we have

$$\sup_{m \in \mathcal{M}_n} \left\{ (2 - \delta_-)p(m) - \text{pen}(m) - \varepsilon \frac{R_{A,m}}{n} \right\}_+ \leq 0.$$

We choose $x = 2\varepsilon$. We obtain that, for Theorem 3.1, the expectation of (5.11) is upper bounded by κn^{-1} and for Theorem 4.1, the term (5.11) is equal to 0 on $\Omega_n \cap \Omega_{\text{pen}}$.

Control of (5.12). Let m, m' in $\mathcal{M}_n$ and let m_s be the index such that $R_{A,m_s} = R_{A,m} \vee R_{A,m'}$ and let κ_1 be a constant to be chosen later. Assume that n is sufficiently large to ensure that $\kappa_1 \varepsilon_n \leq 1/2$. It comes from Fact 6 that

$$\begin{aligned} \delta(m, m') &= f_5((m, m'), \kappa_1) + \kappa_1 \varepsilon_n \frac{R_{A,m_s}}{n} \\ &\leq f_5((m, m'), \kappa_1) + 2\kappa_1 \varepsilon_n \|\widehat{s}_{A,m_s} - s\|^2 + 2\kappa_1 \varepsilon_n \{f_2(m_s, \kappa_1)\}_+. \end{aligned}$$

We deduce from Lemma 5.1 that there exist κ_1 and κ_2 such that

$$\begin{aligned} &\mathbb{E} \left(\sup_{(m, m') \in \mathcal{M}_n^2} \left\{ \delta(m, m') - 2\kappa_1 \varepsilon_n (\|\widehat{s}_{A,m} - s\|^2 + \|\widehat{s}_{A,m'} - s\|^2) \right\} \right) \\ &\leq \mathbb{E} \left(\sup_{(m, m') \in \mathcal{M}_n^2} \left\{ f_5((m, m') + 2\kappa_1 \varepsilon_n f_2(m_s, \kappa_1) \right\}_+ \right) \leq \frac{\kappa_2}{n}. \end{aligned}$$

Applying Lemma 5.2, we obtain a constant κ_1 such that, on Ω_n ,

$$\sup_{(m, m') \in \mathcal{M}_n^2} \left\{ \delta(m, m') - 2\kappa_1 \varepsilon_n (\|\widehat{s}_{A,m} - s\|^2 + \|\widehat{s}_{A,m'} - s\|^2) \right\} \leq 0.$$

Conclusion of the proofs. We use Fact 5 with $x = 2\varepsilon$ and $y = 2\kappa_1 \varepsilon_n$. We take the expectation for the proof of Theorem 3.1, we have obtained that the expectation of the remainder terms (5.10)-(5.12) are upper bounded by κn^{-1} for a sufficiently large n . For the proof of Theorem 4.1, we have obtained that the remainder terms (5.10)-(5.12) with $x = 2\varepsilon$ and $y = 2\kappa_1 \varepsilon_n$ are equal to 0 on $\Omega_n \cap \Omega_{\text{pen}}$ when n is sufficiently large. As explained in the beginning of the proof, this is sufficient to conclude the proof of (3.7) and (4.4).

Let us prove (3.8). Let $\kappa_1 < 1/(2\varepsilon_n)$, from Fact 6 and (3.7), we have

$$\begin{aligned} \frac{K_n}{n} \mathbb{E}(2D_{A,\widehat{m}}) &\leq K_n \left(\mathbb{E}(p(\widehat{m})) + \mathbb{E}(f_2(\widehat{m}, \kappa_1)) + \kappa_1 \varepsilon_n \mathbb{E} \left(\frac{R_{A,\widehat{m}}}{n} \right) \right) \\ &\leq (1 + 2\kappa_1 \varepsilon_n) \mathbb{E}((f_2(\widehat{m}, \kappa_1))_+) + (1 + 2\kappa_1 \varepsilon_n) K_n \mathbb{E}(\|s - \tilde{s}_A\|^2) \\ &\leq 2\mathbb{E}((f_2(\widehat{m}, \kappa_1))_+) + 2K_n \mathbb{E}(\|s - \tilde{s}_A\|^2) \\ &\leq 2 \left(\mathbb{E}((f_2(\widehat{m}, \kappa_1))_+) + \frac{\kappa}{n} \right) + 2R_n. \end{aligned}$$

We used that, by definition $K_n \leq 1$. We conclude the proof with Lemma 5.1. In order to get the bound on $D_{A,\hat{m}}$ in Theorem 4.3, we use that, on $\Omega_n \cap \Omega_{\text{pen}}$, (4.4) holds and there exists a constant κ_1 such that, $\kappa_1 \varepsilon_n < 1/2$ satisfying

$$\begin{aligned} K_n \frac{2D_{A,\hat{m}}}{n} &\leq \frac{K_n}{1 - \kappa_1 \varepsilon_n} (\|s - s_{\hat{m}}\|^2 + p^*(\hat{m})) = \frac{K_n}{1 - \kappa_1 \varepsilon_n} \|s - \tilde{s}_A\|^2 \\ &\leq \frac{1}{1 - \kappa_1 \varepsilon_n} \inf_{m \in \mathcal{M}_n} \|s - \hat{s}_{A,m}\|^2 \leq \frac{1 + \kappa_1 \varepsilon_n}{1 - \kappa_1 \varepsilon_n} \frac{R_n}{n} \leq 3 \frac{R_n}{n}. \end{aligned}$$

6. Appendix. We present in this section some classical collections of models and prove that they satisfy **H4**.

Regular Histograms: Let d be an integer and let S_d be the space of functions t constant on all the intervals $([k/d, (k+1)/d])_{k \in \mathbb{Z}}$. S_d is called the space of regular histograms with size $1/d$. The family $(\psi_k)_{k \in \mathbb{Z}}$, where, for all k in $\mathbb{Z}$, $\psi_k = \sqrt{d} \mathbf{1}_{[k/d, (k+1)/d]}$ is an orthonormal basis of S_d . Let $B_d = \{t \in S_d, \|t\|^2 \leq 1\}$. From Cauchy-Schwarz inequality, we have

$$\sup_{t \in B_d} t^2 = \sum_{k \in \mathbb{Z}} \psi_k^2 = d \mathbf{1}_{\mathbb{R}}.$$

Hence

$$b_m^2 = \left\| \sup_{t \in B_d} t^2 \right\|_{\infty} = d, \quad P \left(\sup_{t \in B_d} t^2 \right) = dP(\mathbf{1}_{\mathbb{R}}) = d.$$

H4 holds on all the spaces S_d , therefore, it holds on the collection $(S_d)_{d=1, \dots, n}$ called the regular histograms collection.

Fourier Spaces: Let $k \geq 1$ be an integer and let, for all x in $[0, 1]$,

$$\psi_{1,k}(x) = \sqrt{2} \cos(2\pi kx), \quad \psi_{2,k}(x) = \sqrt{2} \sin(2\pi kx), \quad \psi_0 = \mathbf{1}_{[0,1]}.$$

Let $\mathcal{M}_n = \{1, \dots, n\}$ and $\forall m \in \mathcal{M}_n$, let $\Lambda_m = \{0, (1, k), (2, k), k = 1, \dots, m\}$. The space S_m , spanned by the family $(\psi_{\lambda})_{\lambda \in \Lambda_m}$ is called the Fourier space with harmonic smaller than m and the collection $(S_m, m \in \mathcal{M}_n)$ is called the collection of Fourier spaces. Let $B_m = \{t \in S_m, \|t\|^2 \leq 1\}$. From Cauchy-Schwarz inequality, for all x in $[0, 1]$,

$$\sup_{t \in B_m} t^2(x) = \sum_{\lambda \in \Lambda_m} \psi_{\lambda}^2(x) = 1 + 2 \sum_{k=1}^m (\cos^2(2\pi kx) + \sin^2(2\pi kx)) = 1 + 2m.$$

Hence, if P is supported in $[0, 1]$,

$$b_m^2 = \left\| \sup_{t \in B_m} t^2 \right\|_{\infty} = 1 + 2m, \quad P \left(\sup_{t \in B_m} t^2 \right) = 1 + 2m.$$

H4 holds on the collection of Fourier spaces when P is supported on $[0, 1]$.

Wavelet Spaces: Assume that $(S_m, m \in \mathcal{M}_n)$ is a collection of wavelet spaces **[W]**. Assume moreover that the scaling function ϕ and the mother wavelet ψ satisfy the following relation. There exists a constant $K_o > 0$ such that, for all x in $\mathbb{R}$,

$$\frac{1}{K_o} \leq \sum_{k \in \mathbb{Z}} \phi^2(x - k) \leq K_o, \quad \frac{1}{K_o} \leq \sum_{k \in \mathbb{Z}} \psi^2(x - k) \leq K_o.$$

This condition is satisfied by the Haar basis, where $\phi = \mathbf{1}_{[0,1]}$, $\psi = \mathbf{1}_{[0,1/2]} - \mathbf{1}_{[1/2,1]}$, with $K_o = 1$. Then, for all $j \geq 0$, we have

$$\frac{1}{K_o} \leq \sum_{k \in \mathbb{Z}} \phi^2(2^j x - k) \leq K_o, \quad \frac{1}{K_o} \leq \sum_{k \in \mathbb{Z}} \psi^2(2^j x - k) \leq K_o.$$

Let $B_m = \left\{ t \in S_m, \|t\|^2 \leq 1 \right\}$. From Cauchy-Schwarz inequality, we have

$$\Psi_m(x) = \sup_{t \in B_m} t^2(x) = \sum_{\lambda \in \Lambda_m} \psi_\lambda^2(x) = \sum_{k \in \mathbb{Z}} 2\phi^2(2x - k) + \sum_{j=1}^{J_m} 2^j \sum_{k \in \mathbb{Z}} \psi^2(2^j x - k).$$

We deduce that

$$\frac{2^{J_m}}{K_o} \leq \frac{1}{K_o} \left(2 + \sum_{j=1}^{J_m} 2^j \right) \leq \Psi_m(x) \leq K_o \left(2 + \sum_{j=1}^{J_m} 2^j \right) \leq 2K_o 2^{J_m}.$$

Hence $b_m^2 = \|\Psi_m\|_\infty \leq 2K_o 2^{J_m}$, $P(\Psi_m) \geq 2^{J_m}/K_o$.

H4 holds on the collection **[W]**.

SUPPLEMENTARY MATERIAL

Supplement A: Proofs of Lemmas 5.1 and 5.2

(<http://lib.stat.cmu.edu/aos/???/???>). In the supplementary material, we give complete proofs of the concentrations lemmas 5.1 and 5.2. We use coupling results respectively of Berbee (1979) and Dedecker and Prieur (2005) to build sequences of independent random variables $(A_0^*, \dots, A_{p-1}^*)$ approximating the sequence of blocks $(A_0, \dots, A_{p-1})$, respectively in the β and τ mixing case. We prove concentration lemmas equivalent to 5.1 and 5.2 for these approximating random variables. The main tools here are the concentration inequalities of Bousquet (2002) and Klein and Rio (2005) for the maximum of the empirical process. We prove finally some covariance inequalities to evaluate the expectation of $p(m)$ and deduce the rates $\varepsilon_n = (\ln n)^{-1/2}$.

References.

- ANDREWS, D. W. K. (1984). Nonstrong mixing autoregressive processes. *J. Appl. Probab.* **21** 930–934. [MR766830 \(86e:60027\)](#)
- ARLOT, S. (2008). *V-fold cross-validation improved: V-fold penalization.* [arXiv:0802.0566v2](#).
- ARLOT, S. (2009). Model Selection by resampling penalization. *Electron. J. Statist.* **3** 557–624.
- ARLOT, S. and MASSART, P. (2009). Data-driven calibration of penalties for least-squares regression. *Journal of Machine learning research* **10** 245–279.
- BARAUD, Y., COMTE, F. and VIENNET, G. (2001). Adaptive estimation in autoregression or β -mixing regression via model selection. *Ann. Statist.* **29** 839–875. [MR1865343 \(2002h:62116\)](#)
- BARRON, A., BIRGÉ, L. and MASSART, P. (1999). Risk bounds for model selection via penalization. *Probab. Theory Related Fields* **113** 301–413. [MR1679028 \(2000k:62049\)](#)
- BERBEE, H. C. P. (1979). *Random walks with stationary increments and renewal theory.* *Mathematical Centre Tracts* **112**. Mathematisch Centrum, Amsterdam. [MR547109 \(81e:60093\)](#)
- BIRGÉ, L. and MASSART, P. (1997). From model selection to adaptive estimation. In *Festschrift for Lucien Le Cam* 55–87. Springer, New York. [MR1462939 \(98m:62086\)](#)
- BIRGÉ, L. and MASSART, P. (2007). Minimal penalties for Gaussian model selection. *Probab. Theory Related Fields* **138** 33–73. [MR2288064 \(2008g:62070\)](#)
- BOUSQUET, O. (2002). A Bennett concentration inequality and its application to suprema of empirical processes. *C. R. Math. Acad. Sci. Paris* **334** 495–500. [MR1890640 \(2003f:60039\)](#)
- BRADLEY, R. C. (2007). *Introduction to strong mixing conditions. Vol. 1.* Kendrick Press, Heber City, UT. [MR2325294](#)
- COMTE, F., DEDECKER, J. and TAUPIN, M. L. (2008). Adaptive density deconvolution with dependent inputs. *Math. Methods Statist.* **17** 87–112. [MR2429122](#)
- COMTE, F. and MERLEVÈDE, F. (2002). Adaptive estimation of the stationary density of discrete and continuous time mixing processes. *ESAIM Probab. Statist.* **6** 211–238 (electronic). New directions in time series analysis (Luminy, 2001). [MR1943148 \(2004b:62105\)](#)
- DEDECKER, J. and PRIEUR, C. (2005). New dependence coefficients. Examples and applications to statistics. *Probab. Theory Related Fields* **132** 203–236. [MR2199291 \(2007b:62081\)](#)
- DEDECKER, J., DOUKHAN, P., LANG, G., LEÓN R., J. R., LOUHICHI, S. and PRIEUR, C. (2007). *Weak dependence: with examples and applications.* *Lecture Notes in Statistics* **190**. Springer, New York. [MR2338725 \(2009a:62009\)](#)
- DONOHU, D. L., JOHNSTONE, I. M., KERKYACHARIAN, G. and PICARD, D. (1996). Density estimation by wavelet thresholding. *Ann. Statist.* **24** 508–539. [MR1394974 \(97f:62061\)](#)
- DOUKHAN, P. (1994). *Mixing. Lecture Notes in Statistics* **85**. Springer-Verlag, New York. Properties and examples. [MR1312160 \(96b:60090\)](#)
- GANNAZ, I. and WINTENBERGER, O. (2009). Adaptive density estimation under dependence. *forthcoming in ESAIM, Probab. and Statist.*
- KLEIN, T. and RIO, E. (2005). Concentration around the mean for maxima of empirical processes. *Ann. Probab.* **33** 1060–1077. [MR2135312 \(2006c:60022\)](#)
- KÜNSCH, H. (1989). The jackknife and the bootstrap for general stationary observations. *Ann. Statist.* **17** 1217–1241.
- LACOUR, C. (2008). Nonparametric estimation of the stationary density and the

- transition density of a Markov chain. *Stochastic Process. Appl.* **118** 232–260. [MR2376901 \(2009b:62073\)](#)
- LERASLE, M. (2009a). Adaptive density estimation of stationary β -mixing and τ -mixing processes. *Math. Methods Statist.* **18** 59–83. [MR2508949 \(2010d:62085\)](#)
- LERASLE, M. (2009b). Optimal model selection in density estimation. *arXiv:0910.1654*.
- LIU, R. and SINGH, K. (1992). Moving block jackknife and bootstrap capture weak dependence. *R. Lepage & L. Billard eds, Exploring the limits of bootstrap (wiley, New York)* 225–248.
- MASSART, P. and NÉDÉLEC, E. (2006). Risk bounds for statistical learning. *Ann. Statist.* **34** 2326–2366. [MR2291502 \(2009e:62282\)](#)
- RUDEMO, M. (1982). Empirical choice of histograms and kernel density estimators. *Scand. J. Statist.* **9** 65–78. [MR668683 \(83j:62059\)](#)
- VOLKONSKIĬ, V. A. and ROZANOV, Y. A. (1959). Some limit theorems for random functions. I. *Teor. Veroyatnost. i Primenen* **4** 186–207. [MR0105741 \(21 ##4477\)](#)

E-MAIL: lerasle@gmail.com

**SUPPLEMENTARY MATERIAL FOR THE PAPER
OPTIMAL MODEL SELECTION FOR DENSITY
ESTIMATION OF STATIONARY DATA UNDER VARIOUS
MIXING CONDITIONS.**

BY MATTHIEU LERASLE*

* *Institut Mathématiques de Toulouse (UMR 5219), INSA Toulouse*

1. Notations. Let $(X_1, \dots, X_n)$ be real valued random variables identically distributed, with common density s with respect to the Lebesgue measure. Let p and q are integers such that $2pq = n$, and that $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$. For all $k = 0, \dots, p-1$, let $I_k = (2kq + 1, \dots, (2k+1)q)$, $A_k = (X_i)_{i \in I_k}$ and $I = \cup_{k=0}^{p-1} I_k$. For all t in $L^2(\mu)$ and all $x_1, \dots, x_q$ in $\mathbb{R}$,

$$L_q(t)(x_1, \dots, x_q) = \frac{1}{q} \sum_{i=1}^q t(x_i), \quad P_A t = \frac{1}{p} \sum_{k=0}^{p-1} L_q(t)(A_k) = \frac{2}{n} \sum_{i \in I} t(X_i),$$

$$\nu_A(t) = (P_A - P)(t).$$

For all m in $\mathcal{M}_n$, we denote by $(\psi_\lambda)_{\lambda \in \Lambda_m}$ an orthonormal basis of S_m . The estimator $\hat{s}_{A,m}$ associated to the model S_m , is defined as

$$\hat{s}_{A,m} = \sum_{\lambda \in \Lambda_m} (P_A \psi_\lambda) \psi_\lambda.$$

Classical computations show that, if s_m denotes the orthogonal projection of s onto S_m ,

$$s_m = \sum_{\lambda \in \Lambda_m} (P \psi_\lambda) \psi_\lambda, \quad \text{hence } \|\hat{s}_{A,m} - s_m\|^2 = \sum_{\lambda \in \Lambda_m} (\nu_A \psi_\lambda)^2.$$

For all m, m' in $\mathcal{M}_n$, let

$$p(m) = \|s_m - \hat{s}_{A,m}\|^2 = \sup_{t \in B_m} (\nu_A(t))^2 = \sum_{\lambda \in \Lambda_m} (\nu_A(\psi_\lambda))^2.$$

AMS 2000 subject classifications: Primary 62G07, 62G09; secondary 62M99

Keywords and phrases: Density estimation, optimal model selection, resampling methods, slope heuristic, weak dependence

$$\delta(m, m') = 2\nu_A(s_m - s_{m'}).$$

Hereafter $W_0, \dots, W_{p-1}$ denotes a resampling scheme, $\overline{W} = p^{-1} \sum_{i=0}^{p-1} W_i$, P_A^W denotes the resampling empirical process, defined for all measurable functions t by

$$P_A^W t = \frac{1}{p} \sum_{i=0}^{p-1} W_i L_q t(A_i).$$

We introduce also $\nu_A^W = P_A^W - \overline{W} P_A$ and $\tilde{C}_W = (\text{Var}(W_1 - \overline{W}))^{-1}$. For any orthonormal basis $(\psi_\lambda)_{\lambda \in \Lambda_m}$ of S_m , let

$$p_W(m) = \tilde{C}_W \sum_{\lambda \in \Lambda_m} \mathbb{E}_W ((\nu_A^W(\psi_\lambda))^2).$$

Let $\varepsilon_n = (\ln n)^{-1/2}$ and let $\kappa > 0$. Let $\mathcal{M}$ denote one of the set $\mathcal{M}_n$ or $\mathcal{M}_n^2$. When $\mathcal{M} = \mathcal{M}_n$, for all $\overline{m}$ in $\mathcal{M}$ let $R_{A,\overline{m}} = R_{A,m}$ and when $\mathcal{M} = \mathcal{M}_n^2$, for all $\overline{m} = (m, m')$ in $\mathcal{M}$, let $R_{A,\overline{m}} = R_{A,m} \vee R_{A,m'}$. For all $\overline{m}$ in $\mathcal{M}$, let

$$(1.1) \quad f_1(\overline{m}, \kappa) = p(m) - \frac{2D_{A,m}}{n} - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(1.2) \quad f_2(\overline{m}, \kappa) = \frac{2D_{A,m}}{n} - p(m) - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(1.3) \quad f_3(\overline{m}, \kappa) = p(m) - p_W(m) - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(1.4) \quad f_4(\overline{m}, \kappa) = p_W(m) - p(m) - \kappa \varepsilon_n \frac{R_{A,m}}{n},$$

$$(1.5) \quad f_5(\overline{m}, \kappa) = \delta(m, m') - \kappa \varepsilon_n \frac{R_{A,m} \vee R_{A,m'}}{n}.$$

In this paper, we state and prove the following results.

LEMMA 1.1. *Let $X_1, \dots, X_n$ be a strictly stationary sequence of real valued random variables with common density s and let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of regular wavelet spaces **[W]** satisfying assumptions **H3, H4**. Let p, q be two integers satisfying $2pq = n$ and $\frac{1}{2}\sqrt{n}(\ln n)^2 \leq p \leq \sqrt{n}(\ln n)^2$. Assume that there exists $\theta > 5$ such that $X_1, \dots, X_n$ are arithmetically **[AR](θ)** τ -mixing and satisfy **S**($\tau, \mathbf{W}$). There exist constants κ_1, κ_2 , such that, for all $i = 1, \dots, 5$, for all $\overline{m}$ in $\mathcal{M}$,*

$$(1.6) \quad \mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} (f_i(\overline{m}, \kappa_1))_+ \right) \leq \frac{\kappa_2}{n}.$$

LEMMA 1.2. *Let $\theta > 1$ and let $(X_n)_{n \in \mathbb{Z}}$ be an arithmetically $[\mathbf{AR}(\theta)]$ β -mixing process satisfying $\mathbf{S}(\beta)$. Let $(S_m)_{m \in \mathcal{M}_n}$ be a collection of linear spaces satisfying Assumptions **H1**, **H2**, **H3**, **H4**. Let p, q such that $2pq = n$, $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$. There exist constants κ_1, κ_2 which may vary from line to line such that, on an event Ω_n satisfying*

$$\mathbb{P}(\Omega_n) \geq 1 - \kappa_2 \left(\frac{(\ln n)^{2(1+\theta)}}{n^{\theta/2}} \vee \frac{1}{n^2} \right), \text{ where}$$

$$(1.7) \quad \forall \overline{m} \in \mathcal{M}, \forall i = 1, \dots, 5, f_i(\overline{m}, \kappa_1) \leq 0.$$

This paper is not self-contained. Some notations and assumptions have been introduced in the main paper. These lemmas are central in the proof of the oracle inequalities in the main paper. We believe that the tools involved are interesting by themselves and we group them in three subsections. The first one gives technical results on the resampling penalty. The second gives coupling results for mixing data and the third one provides the useful concentration inequalities for independent processes. The proof of the Lemmas are given respectively in Sections 6.1 and 6.2.

2. Some results on resampling penalties.

LEMMA 2.1. *Let $(t_\lambda)_{\lambda \in \Lambda}$ be a set of real valued square integrable functions defined on a measurable space $(\mathbb{X}, \mathcal{X})$. Let $A_0, \dots, A_{p-1}$ be $\mathbb{X}$ -valued random variables with common law P and let $(W_0, \dots, W_{p-1})$ be a resampling scheme of $A_0, \dots, A_{p-1}$. for all t , let*

$$P_A t = \frac{1}{p} \sum_{i=0}^{p-1} t(A_i), \quad \nu_A t = (P_A - P)t, \quad p(\Lambda) = \sum_{\lambda \in \Lambda} (\nu_A(t_\lambda))^2.$$

Let $\overline{W} = p^{-1} \sum_{i=0}^{p-1} W_i$ and $\tilde{C}_W = \text{Var}(W_1 - \overline{W})^{-1}$. Let

$$\nu_A^W t = \frac{1}{p} \sum_{i=0}^{p-1} (W_i - \overline{W}) t(A_i), \quad p_W(\Lambda) = \tilde{C}_W \sum_{\lambda \in \Lambda} E_W ((\nu_A^W(t_\lambda))^2),$$

$T(\Lambda) = \sum_{\lambda \in \Lambda} (t_\lambda - P t_\lambda)^2$, $D = PT$ and

$$U = \frac{1}{p(p-1)} \sum_{i \neq j=0}^{p-1} \sum_{\lambda \in \Lambda} (t_\lambda(A_i) - t_\lambda)(t_\lambda(A_j) - P t_\lambda).$$

Then, we have

$$(2.1) \quad p(\Lambda) = \frac{1}{p}P_AT + \frac{p-1}{p}U, \quad p_W(\Lambda) = \frac{1}{p}(P_AT - U), \quad p(\Lambda) - p_W(\Lambda) = U.$$

When, moreover, $A_0, \dots, A_{p-1}$ are i.i.d,

$$(2.2) \quad \mathbb{E}(p_W(\Lambda)) = \frac{D}{p}, \quad p_W(\Lambda) - \frac{D}{p} = \frac{1}{p}(\nu_AT - U).$$

Proof: In the independent case, it is clear that $\mathbb{E}(P_AT) = D$ and $\mathbb{E}(U) = 0$, thus (2.2) comes from (2.1). We only have to prove the two first equalities, the third one being an immediate consequence.

$$\begin{aligned} p(\Lambda) &= \frac{1}{p^2} \sum_{i=0}^{p-1} \sum_{\lambda \in \Lambda} (t_\lambda(A_i) - Pt_\lambda)^2 \\ &\quad + \frac{1}{p^2} \sum_{i \neq j=0}^{p-1} \sum_{\lambda \in \Lambda} (t_\lambda(A_i) - Pt_\lambda)(t_\lambda(A_j) - Pt_\lambda) = \frac{1}{p}P_AT + \frac{p-1}{p}U. \end{aligned}$$

Since $\sum_i (W_i - \overline{W}) = 0$, we have

$$(2.3) \quad \nu_A^W(t_\lambda) = \frac{1}{p} \sum_{i=0}^{p-1} (W_i - \overline{W})t_\lambda(A_i) = \frac{1}{p} \sum_{i=0}^{p-1} (W_i - \overline{W})(t_\lambda(A_i) - Pt_\lambda).$$

Let $E_{i,j} = \tilde{C}_W \mathbb{E}[(W_i - \overline{W})(W_j - \overline{W})]$. Since the weights are exchangeables, for all $i = 1, \dots, n$, $E_{i,i} = E_{1,1} = 1$ and for all $i \neq j$, $E_{i,j} = E_{1,2}$. Taking the square in (2.3), we obtain that

$$\begin{aligned} \tilde{C}_W \mathbb{E}_W(\nu_A^W(t_\lambda))^2 &= \frac{1}{p^2} \sum_{i,j=0}^{p-1} E_{i,j} (t_\lambda(A_i) - Pt_\lambda)(t_\lambda(A_j) - Pt_\lambda) \\ &= \frac{1}{p^2} \sum_{i=0}^{p-1} (t_\lambda(A_i) - Pt_\lambda)^2 + \frac{E_{1,2}}{p^2} \sum_{i \neq j=0}^{p-1} (t_\lambda(A_i) - Pt_\lambda)(t_\lambda(A_j) - Pt_\lambda). \end{aligned}$$

Summing in λ , we obtain that

$$p_W(\Lambda) = \frac{1}{p}P_AT + \frac{p-1}{p}E_{1,2}U.$$

Since $\sum_i (W_i - \overline{W}) = 0$,

$$0 = \mathbb{E} \left[\left(\sum_{i=0}^{p-1} (W_i - \overline{W}) \right)^2 \right] = \tilde{C}_W^{-1} (p + p(p-1)E_{1,2}).$$

Hence $E_{1,2} = -(p-1)^{-1}$, and the proof is concluded.

3. Coupling results. The first coupling lemma was obtained in [Lerasle \(2009a\)](#), as a consequence of a result from [Dedecker and Prieur \(2005\)](#).

LEMMA 3.1. [τ -coupling, Claim 1 p17 in [Lerasle \(2009a\)](#)] Assume that the process $(X_1, \dots, X_n)$ is τ -mixing and let p, q and $A_0, \dots, A_{p-1}$ be respectively the integers and the random variables defined in Section 1. There exist random variables $A_0^*, \dots, A_{p-1}^*$ such that:

1. for all $k = 0, \dots, p-1$, $A_k^* = (X_{2kq+1}^*, \dots, X_{(2k+1)q}^*)$ has the same law as A_k ,
2. for all $k = 0, \dots, p-1$, A_k^* is independent of $A_0, \dots, A_{k-1}, A_0^*, \dots, A_{k-1}^*$,
3. for all $k = 0, \dots, p-1$, $\mathbb{E}(d_q(A_k, A_k^*)) \leq q\tau_q$.

β -mixing data satisfy the very important following lemma, which is due to [Viennet \(1997\)](#).

LEMMA 3.2. (Lemma 5.1 in [Viennet \(1997\)](#)) Assume that the process $(X_1, \dots, X_n)$ is β -mixing and let p, q and $A_0, \dots, A_{p-1}$ be respectively the integers and the random variables defined in Section 1. There exist random variables $A_0^*, \dots, A_{p-1}^*$ such that:

1. for all $k = 0, \dots, p-1$, $A_k^* = (X_{2kq+1}^*, \dots, X_{(2k+1)q}^*)$ has the same law as A_k ,
2. for all $k = 0, \dots, p-1$, A_k^* is independent of $A_0, \dots, A_{k-1}, A_0^*, \dots, A_{k-1}^*$,
3. for all $k = 0, \dots, p-1$, $\mathbb{P}(A_k \neq A_k^*) \leq \beta_q$.

For all functionals $T = F(A_0, \dots, A_{p-1})$, let $T^* = F(A_0^*, \dots, A_{p-1}^*)$, where the random variables (A_k^*) are given by the previous coupling lemmas. In particular, we will use repeatedly the notations $P_A^* = \frac{2}{n} \sum_{i \in I} \delta_{X_i^*}$, $\nu_A^* = P_A^* - P$, $(P_A^W)^* = \frac{2}{n} \sum_{k=0}^{p-1} W_k \sum_{i \in I_k} \delta_{X_i^*}$, $(\nu_A^W)^* = (P_A^W)^* - \bar{W}P_A^*$, $\delta^*(m, m') = 2\nu_A^*(s_m - s_{m'})$,

$$(3.1) \quad p^*(m) = \sum_{\lambda \in \Lambda_m} (\nu_A^* \psi_\lambda)^2, p_W^*(m) = \tilde{C}_W \sum_{\lambda \in \Lambda_m} ((\nu_A^W)^* \psi_\lambda)^2.$$

$$U_m = \frac{1}{p(p-1)} \sum_{\lambda \in \Lambda_m} \sum_{i \neq j=0}^{p-1} (L_q(\psi_\lambda)(A_i) - P\psi_\lambda)(L_q(\psi_\lambda)(A_j) - P\psi_\lambda)$$

$$U_m^* = \frac{1}{p(p-1)} \sum_{\lambda \in \Lambda_m} \sum_{i \neq j=0}^{p-1} (L_q(\psi_\lambda)(A_i^*) - P\psi_\lambda)(L_q(\psi_\lambda)(A_j^*) - P\psi_\lambda)$$

The first Lemma is a straightforward consequence of Lemma 3.2.

LEMMA 3.3. *Let $X_1, \dots, X_n$ be real valued, stationary, β -mixing random variables. Let p, q be two integers such that $2pq = n$ and let $(S_m)_{m \in \mathcal{M}_n}$ be any collection of spaces of measurable functions. Let $A_0^*, \dots, A_{p-1}^*$ be the independent random variables given by Lemma 3.2. Let $p, p_W, \delta, p^*, p_W^*$ and δ^* be the associated functions defined on $\mathcal{M}_n$ and $\mathcal{M}_n^2$ in (3.1). There exists an event $\Omega_n^{(1)}$ with $\mathbb{P}(\Omega_n^{(1)}) \geq 1 - p\beta_q$ where, for all m, m' in $\mathcal{M}_n$,*

$$p(m) = p^*(m), p_W(m) = p_W^*(m), \delta(m, m') = \delta^*(m, m'), U_m = U_m^*.$$

Proof: Consider the event $\Omega_n^{(1)} = \{\forall l = 0, \dots, p-1, A_l = A_l^*\}$. It comes from Viennet's coupling lemma that $\mathbb{P}(\Omega_n^{(1)}) \geq 1 - p\beta_q$ and it is clear that, on $\Omega_n^{(1)}$, the conclusion of Lemma 3.3 holds.

LEMMA 3.4. *Let $X_1, \dots, X_n$ be stationary random variables, real valued, τ -mixing and with common density s . Let p and q be two integers such that $2pq = n$ and let $A_0^*, \dots, A_{p-1}^*$ be the random variables given by Lemma 3.1. Let $\mathcal{M}_n$ be a collection of models. Let $p, p_W, \delta, p^*, p_W^*, \delta^*$ be the associated functions defined on $\mathcal{M}_n$ and $(\mathcal{M}_n)^2$ in (3.1). Let MC_n be the mixing complexity of $\mathcal{M}_n$ defined by*

$$MC_n = \sum_{m \in \mathcal{M}_n} \left(\left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda) + \|s\| |\mathcal{M}_n| \sup_{t \in B_m} \text{Lip}(t) \right).$$

Then

$$(3.2) \quad \mathbb{E} \left(\sup_{m \in \mathcal{M}_n} |p(m) - p^*(m)| \right) \leq 4\tau_q MC_n,$$

$$(3.3) \quad \mathbb{E} \left(\sup_{m \in \mathcal{M}_n} |p_W(m) - p_W^*(m)| \right) \leq \frac{8\tau_q}{p} MC_n,$$

$$(3.4) \quad \mathbb{E} \left(\sup_{m, m' \in \mathcal{M}_n} |\delta(m, m') - \delta^*(m, m')| \right) \leq 4\tau_q MC_n.$$

Proof: For all m in $\mathcal{M}_n$, we have

$$\mathbb{E} \left(\sup_{m \in \mathcal{M}_n} |p(m) - p^*(m)| \right) \leq \sum_{m \in \mathcal{M}_n} \mathbb{E} (|p(m) - p^*(m)|).$$

Moreover, for all m in $\mathcal{M}_n$,

$$\begin{aligned}
|p(m) - p^*(m)| &= \left| \sum_{\lambda \in \Lambda_m} ((P_A - P)\psi_\lambda)^2 - ((P_A^* - P)\psi_\lambda)^2 \right| \\
&= \left| \sum_{\lambda \in \Lambda_m} ((\nu_A + \nu_A^*)\psi_\lambda) ((P_A - P_A^*)\psi_\lambda) \right| \\
|p(m) - p^*(m)| &\leq \sum_{\lambda \in \Lambda_m} |(\nu_A + \nu_A^*)\psi_\lambda| \frac{1}{p} \sum_{k=0}^{p-1} |L_q(\psi_\lambda)(A_k) - L_q(\psi_\lambda)(A_k^*)| \\
&\leq 4 \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}_q(L_q(\psi_\lambda)) \frac{1}{p} \sum_{k=0}^{p-1} d_q(A_k, A_k^*) \\
&\leq \frac{4}{q} \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda) \frac{1}{p} \sum_{k=0}^{p-1} d_q(A_k, A_k^*).
\end{aligned}$$

We take the expectation in this last inequality and we use Lemma 3.1 to obtain (3.2). From Lemma 2.1,

$$|p_W(m) - p_W^*(m)| = \frac{1}{p} |(P_A - P_A^*)(T_m) - (U_m - U_m^*)|.$$

$(P_A - P_A^*)T_m$ is equal to

$$\sum_{\lambda \in \Lambda_m} \frac{1}{p} \sum_{k=0}^{p-1} (L_q(\psi_\lambda)(A_k) - L_q(\psi_\lambda)(A_k^*)) (L_q(\psi_\lambda)(A_k) + L_q(\psi_\lambda)(A_k^*) - 2P\psi_\lambda),$$

thus

$$\begin{aligned}
|(P_A - P_A^*)T_m| &= 4 \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}_q(L_q(\psi_\lambda)) \frac{1}{p} \sum_{k=0}^{p-1} d_q(A_k, A_k^*) \\
&\leq \frac{4}{q} \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda) \frac{1}{p} \sum_{k=0}^{p-1} d_q(A_k, A_k^*).
\end{aligned}$$

Moreover

$$\begin{aligned}
U_m - U_m^* &= \frac{1}{p(p-1)} \sum_{i \neq j=0}^{p-1} \sum_{\lambda \in \Lambda_m} (L_q(\psi_\lambda)(A_j) - P\psi_\lambda)(L_q(\psi_\lambda)(A_i) - L_q(\psi_\lambda)(A_i^*)) \\
&\quad + \frac{1}{p(p-1)} \sum_{i \neq j=0}^{p-1} \sum_{\lambda \in \Lambda_m} (L_q(\psi_\lambda)(A_i^*) - P\psi_\lambda)(L_q(\psi_\lambda)(A_j) - L_q(\psi_\lambda)(A_j^*)),
\end{aligned}$$

thus

$$|U_m - U_m^*| \leq \frac{4}{q} \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda) \frac{1}{p} \sum_{k=0}^{p-1} d_q(A_k, A_k^*).$$

Therefore,

$$\begin{aligned} \mathbb{E}(|p_W(m) - p_W^*(m)|) &\leq \frac{8}{pq} \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda) \frac{1}{p} \sum_{k=0}^{p-1} \mathbb{E}(d_q(A_k, A_k^*)) \\ &\leq \frac{8\tau_q}{p} \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda). \end{aligned}$$

Thus $\mathbb{E}(\sup_{m \in \mathcal{M}_n} |p_W(m) - p_W^*(m)|)$ is upper bounded by

$$\sum_{m \in \mathcal{M}_n} \mathbb{E}(|p_W(m) - p_W^*(m)|) \leq \frac{8\tau_q}{p} \sum_{m \in \mathcal{M}_n} \left\| \sum_{\lambda \in \Lambda_m} |\psi_\lambda| \right\|_\infty \sup_{\lambda \in \Lambda_m} \text{Lip}(\psi_\lambda).$$

$\mathbb{E}(\sup_{m, m' \in \mathcal{M}_n} \delta(m, m') - \delta^*(m, m')) \leq \sum_{m, m' \in \mathcal{M}_n} \mathbb{E}(|\delta(m, m') - \delta^*(m, m')|)$
and, for all m, m' in $\mathcal{M}_n$,

$$\begin{aligned} \mathbb{E}(|\delta(m, m') - \delta^*(m, m')|) &= 2\mathbb{E}(|(P_A - P_A^*)(s_m - s_{m'})|) \\ &\leq \frac{2}{pq} \text{Lip}(s_m - s_{m'}) \sum_{k=0}^{p-1} \mathbb{E}(d_q(A_k, A_k^*)) \leq 2\tau_q \text{Lip}(s_m - s_{m'}). \end{aligned}$$

For all x, y in $\mathbb{R}$ and all m, m' in $\mathcal{M}_n$,

$$(s_m - s_{m'})(x) - (s_m - s_{m'})(y) \leq \|s\| \left(\sup_{t \in B_m} \text{Lip}(t) + \sup_{t \in B_{m'}} \text{Lip}(t) \right) |x - y|.$$

Hence, $\text{Lip}(s_m - s_{m'}) \leq \|s\| \left(\sup_{t \in B_m} \text{Lip}(t) + \sup_{t \in B_{m'}} \text{Lip}(t) \right)$, thus

$$\mathbb{E} \left(\sup_{m, m' \in \mathcal{M}_n} \delta(m, m') - \delta^*(m, m') \right) \leq 4\tau_q \|s\| |\mathcal{M}_n| \sum_{m \in \mathcal{M}_n} \sup_{t \in B_m} \text{Lip}(t).$$

LEMMA 3.5. *Assume that $2pq = n$, that $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$, and that there exist constants C and $\theta > 0$ such that, for all q in $\mathbb{N}^*$, $\beta_q \leq Cq^{-(1+\theta)}$. Then*

$$p\beta_q \leq \frac{(\ln n)^{4+2\theta}}{n^{\theta/2}}.$$

Proof: The proof is straightforward.

LEMMA 3.6. *Assume that there exist constants $\kappa_- > 0$, $\kappa_+ > 0$ such that $\kappa_- \sqrt{n}(\ln n)^{-2} \leq q \leq \kappa_+ \sqrt{n}(\ln n)^{-2}$ and assume that, for all q , $\tau_q \leq q^{-(1+\theta)}$. Let MC_n be the mixing complexity defined in Lemma 3.4 for the collection of models $[\mathbf{W}]$. Then, there exists a constant κ such that*

$$\tau_q MC_n \leq \kappa \frac{(\ln n)^{2(1+\theta)}}{n^{(\theta-3)/2}}.$$

Proof: Let us first recall some basic inequalities that hold in $[\mathbf{W}]$: let $K_\infty = (\sqrt{2}\|\phi\|_\infty) \vee \|\psi\|_\infty$, $K_L = (2\sqrt{2}\text{Lip}(\phi)) \vee \text{Lip}(\psi)$, $K_{BV} = AK_L$. Then for all $j \geq 0$, we have $\|\psi_{j,k}\|_\infty \leq K_\infty 2^{j/2}$,

$$(3.5) \quad \left\| \sum_{k \in \mathbb{Z}} |\psi_{j,k}| \right\|_\infty \leq AK_\infty 2^{j/2},$$

$$(3.6) \quad \text{Lip}(\psi_{j,k}) \leq K_L 2^{3j/2},$$

$$(3.7) \quad \|\psi_{j,k}\|_{BV} \leq K_{BV} 2^{j/2}.$$

Since $\text{Card}(\mathcal{M}_n) \leq \ln n / \ln 2$, we obtain that

$$MC_n \leq \kappa \left(\sum_{j=0}^{\lfloor \ln n \rfloor / \ln 2} 2^{2j} + (\ln n) 2^{3j/2} \right) \leq \kappa n^2.$$

We conclude the proof of Lemma 3.6, saying that $q \geq \kappa_- \sqrt{n}(\ln n)^{-2}$ implies

$$\tau_q \leq \frac{C}{\kappa_-^{1+\theta}} \frac{(\ln n)^{2(1+\theta)}}{n^{(1+\theta)/2}}.$$

4. Concentration inequalities. The following concentration inequalities were proved in Lerasle (2009b). These inequalities derive Bousquet's and Klein & Rio's versions of Talagrand's concentration inequality for the supremum of the empirical process (see Bousquet (2002); Klein and Rio (2005)).

THEOREM 4.1. *Let $A_0, \dots, A_{p-1}$ be iid random variables valued in a measurable space $(\mathbb{X}, \mathcal{X})$, with common law P . Let S be a symmetric class of functions bounded by b . For all t in S , let $P_A t = p^{-1} \sum_{i=0}^{p-1} t(A_i)$, $v^2 = \sup_{t \in S} P[(t - Pt)^2]$, $Z = \sup_{t \in S} (P_A - P)t$, $D = p\mathbb{E}(Z^2)$. For all $x > 0$ and all ϵ in $(0, 1]$, with probability larger than $1 - 2e^{-x}$,*

$$\left| Z^2 - \frac{D}{p} \right| \leq \kappa \left(\epsilon \frac{D}{p} + \frac{1}{\epsilon} \left(\frac{v^2 x}{p} + \left(\frac{bx}{\epsilon p} \right)^2 \right) \right).$$

The constant $\kappa = (16(\ln 2)^{-2} + 8)$ works.

Proof: We repeat the proof of this theorem for the sake of completeness. From Bousquet's and Klein & Rio's bound, we have, for all $x > 0$, for all $0 < \epsilon \leq 1$,

$$(4.1) \quad P \left(|Z - \mathbb{E}(Z)| > \epsilon \mathbb{E}(Z) + \sqrt{\frac{2v^2 x}{p}} + 2\epsilon^{-1} \frac{bx}{p} \right) \leq 2e^{-x}.$$

Let then $\Omega^x = \left\{ |Z - \mathbb{E}(Z)| \leq \epsilon \mathbb{E}(Z) + \sqrt{\frac{2v^2 x}{p}} + 2\epsilon^{-1} \frac{bx}{p} \right\}$. On Ω^x ,

$$Z^2 \leq (1 + \epsilon)^3 (\mathbb{E}(Z))^2 + 8\epsilon^{-1} \frac{v^2 x}{p} + 16\epsilon^{-3} \frac{b^2 x^2}{p^2}.$$

Using Cauchy-Schwarz inequality, we have $(\mathbb{E}(Z))^2 \leq \mathbb{E}(Z^2) = p^{-1}D$, since $\epsilon \leq 1$, $(1 + \epsilon)^3 \leq 1 + 7\epsilon$, thus, on Ω^x ,

$$Z^2 \leq (1 + 7\epsilon) \frac{D}{p} + 8\epsilon^{-1} \frac{v^2 x}{p} + 16\epsilon^{-3} \frac{b^2 x^2}{p^2}.$$

As S is symmetric, Z is positive. Hence, we also have, on Ω^x ,

$$\begin{aligned} Z^2 &\geq (1 - \epsilon)^2 (\mathbb{E}(Z))^2 - 2(1 - \epsilon) \mathbb{E}(Z) \left(\sqrt{\frac{2v^2 x}{p}} + 2\epsilon^{-1} \frac{bx}{p} \right) \\ &\geq (1 - 4\epsilon) (\mathbb{E}(Z))^2 - \frac{4}{\epsilon} \frac{v^2 x}{p} - \frac{8}{\epsilon^3} \frac{b^2 x^2}{p^2}. \end{aligned}$$

From Lemma 4.2 above, we have, for $\kappa = 2 + 5\pi/2$,

$$(\mathbb{E}(Z))^2 \geq (1 - 3\epsilon^2) \mathbb{E}(Z^2) - \kappa \frac{v^2}{p} - \frac{16}{\epsilon^2} \frac{b^2}{p^2}.$$

Hence, on Ω^x ,

$$Z^2 \geq (1 - 19\epsilon)\mathbb{E}(Z^2) - \left(\kappa + \frac{4x}{\epsilon}\right) \frac{v^2}{p} - \left(\frac{16}{\epsilon^2} + \frac{8x^2}{\epsilon^3}\right) \frac{b^2}{p^2}.$$

Let us say that we can take $x \geq \ln 2$, otherwise, the conclusion of the theorem is trivial. Moreover, as $\epsilon \leq 1$, we have $1 \leq ((\ln 2)\epsilon)^{-1}x$, hence

$$Z^2 \leq (1 - 24\epsilon)\mathbb{E}(Z^2) - \left(4 + \frac{\kappa}{\ln 2}\right) \frac{v^2 x}{\epsilon p} - \left(\frac{16}{(\ln 2)^2} + 8\right) \frac{b^2}{\epsilon^3 p^2}.$$

Let us now prove Lemma 4.2.

LEMMA 4.2. *Let $A_0, \dots, A_{p-1}$ be iid random variables valued in a measurable space $(\mathbb{X}, \mathcal{X})$, with common law P . Let S be a symmetric class of functions bounded by b . For all t in S , let $P_A t = p^{-1} \sum_{i=0}^{p-1} t(A_i)$, $v^2 = \sup_{t \in S} P[(t - Pt)^2]$, $Z = \sup_{t \in S} (P_A - P)t$. For all $\epsilon > 0$, we have*

$$\text{Var}(Z) \leq 3(\epsilon \mathbb{E}(Z))^2 + \left(2 + \frac{5\pi}{2}\right) \frac{v^2}{p} + \frac{16}{\epsilon^2} \frac{b^2}{p^2}.$$

Proof of Lemma 4.2: We have

$$\begin{aligned} \text{Var}(Z) &= \mathbb{E}(|Z - \mathbb{E}(Z)|^2) = \int_0^\infty P(|Z - \mathbb{E}(Z)|^2 > x) dx \\ &= \int_0^\infty P(|Z - \mathbb{E}(Z)| > x) 2x dx. \end{aligned}$$

Let now ϵ in $(0, 1]$, we have, using the following change of variables $x = \epsilon \mathbb{E}(Z) + \sqrt{\frac{2v^2 y}{p}} + 2\epsilon^{-1} \frac{by}{p}$ and the concentration inequality (4.1),

$$\begin{aligned} \text{Var}(Z) &= \int_0^{\epsilon \mathbb{E}(Z)} P(|Z - \mathbb{E}(Z)| > x) 2x dx \\ &\quad + \int_{\epsilon \mathbb{E}(Z)}^\infty P(|Z - \mathbb{E}(Z)| > x) 2x dx \leq (\epsilon \mathbb{E}(Z))^2 \\ &\quad + 2 \int_0^\infty e^{-y} \left(\epsilon \mathbb{E}(Z) + \sqrt{\frac{2v^2 y}{p}} + 2\epsilon^{-1} \frac{by}{p} \right) \left(\sqrt{\frac{v^2}{2py}} + 2\epsilon^{-1} \frac{b}{p} \right) dy \\ &= (\epsilon \mathbb{E}(Z))^2 + \epsilon \mathbb{E}(Z) \sqrt{2 \frac{v^2}{p}} \int_0^\infty \frac{e^{-y}}{\sqrt{y}} dy + 4\mathbb{E}(Z) \frac{b}{p} \int_0^\infty e^{-y} dy \\ &\quad + 2 \frac{v^2}{p} \int_0^\infty e^{-y} dy + \frac{4}{\epsilon} \sqrt{\frac{v^2}{p}} \frac{b}{p} \int_0^\infty \left(\sqrt{2y} + \frac{1}{\sqrt{2y}} \right) e^{-y} dy \\ &\quad + \frac{8}{\epsilon^2} \frac{b^2}{p^2} \int_0^\infty y e^{-y} dy. \end{aligned}$$

Using that $\int_0^\infty e^{-y} dy = \int_0^\infty y e^{-y} dy = 1$ and

$$\int_0^\infty \frac{e^{-y}}{\sqrt{y}} dy = 2 \int_0^\infty \sqrt{y} e^{-y} dy = \sqrt{\pi},$$

we obtain, using repeatedly the inequality $2ab \leq a^2 + b^2$,

$$\begin{aligned} \text{Var}(Z) &\leq (\epsilon \mathbb{E}(Z))^2 + 2\epsilon \mathbb{E}(Z) \sqrt{\frac{\pi v^2}{2p}} + 4\mathbb{E}(Z) \frac{b}{p} + 2\frac{v^2}{p} + \frac{8}{\epsilon} \sqrt{\frac{\pi v^2}{2p}} \frac{b}{p} + \frac{8}{\epsilon^2} \frac{b^2}{p^2} \\ &\leq 3(\epsilon \mathbb{E}(Z))^2 + \left(2 + \frac{5\pi}{2}\right) \frac{v^2}{p} + \frac{16}{\epsilon^2} \frac{b^2}{p^2}. \square \end{aligned}$$

We deduce from this theorem the following concentration inequality for U -statistics.

COROLLARY 4.3. *Let $A_0, \dots, A_{p-1}$ be i.i.d random variables valued in a measurable space $(\mathbb{X}, \mathcal{X})$, with common law P . Let μ be a measure on $(\mathbb{X}, \mathcal{X})$ and let $(t_\lambda)_{\lambda \in \Lambda_m}$ be a set of functions in $L^2(\mu)$. Let*

$$B = \left\{ t = \sum_{\lambda \in \Lambda_m} a_\lambda t_\lambda, \sum_{\lambda \in \Lambda_m} a_\lambda^2 \leq 1 \right\}, \quad D = \mathbb{E} \left(\sup_{t \in B} (t(A_1) - Pt)^2 \right),$$

$$v^2 = \sup_{t \in B} P[(t - Pt)^2], \quad b = \sup_{t \in B} \|t\|_\infty.$$

Let U be the following U -statistics

$$U = \frac{1}{p(p-1)} \sum_{i \neq j=0}^{p-1} (t_\lambda(A_i) - Pt_\lambda)(t_\lambda(A_j) - Pt_\lambda).$$

For all $x > 0$, with probability larger than $1 - 4e^{-x}$,

$$|U| \leq \kappa' \left(\epsilon \frac{D}{p} + \frac{1}{\epsilon} \left(\frac{v^2 x}{p} + \left(\frac{bx}{\epsilon p} \right)^2 \right) \right),$$

where the constant κ' can be taken equal to $2(\kappa + 4(\ln 2)^{-1})$, for the constant κ defined in Theorem 4.1.

Proof of Corollary 4.3. From Lemma 2.1, we have $U = p(p-1)^{-1}(Z^2 - p^{-1}P_A T)$, where

$$Z = \sqrt{\sum_{\lambda \in \Lambda_m} ((P_A - P)t_\lambda)^2} = \sup_{t \in B} (P_A - P)t, \quad T = \sum_{\lambda \in \Lambda_m} (t_\lambda - Pt_\lambda)^2 = \sup_{t \in B} (t - Pt)^2.$$

From the same lemma, we also have $\mathbb{E}(Z^2) = p^{-1}PT$, so that

$$(4.2) \quad U = p(p-1)^{-1} \left(Z^2 - \mathbb{E}(Z^2) - \frac{1}{p}(P_A - P)T \right).$$

From Bernstein's inequality, we have, with probability larger than $1 - 2e^{-x}$,

$$|(P_A - P)T| \leq \sqrt{\frac{2\text{Var}(T(A_1))x}{p}} + \frac{\|T\|_\infty x}{3p}.$$

We have

$$\|T\|_\infty = \left\| \sup_{t \in B} (t - Pt) \right\|^2 \leq 4b^2.$$

Moreover

$$\text{Var}(T(A_1)) \leq \|T\|_\infty PT \leq 4b^2 D.$$

Hence, with probability $1 - 2e^{-x}$,

$$(4.3) \quad |(P_A - P)T| \leq \sqrt{\frac{8b^2 Dx}{p}} + \frac{4b^2 x}{3p} \leq \epsilon D + \left(\frac{2}{\epsilon} + \frac{4}{3} \right) \frac{b^2 x}{p} \leq \epsilon D + \frac{4}{(\ln 2)\epsilon} \frac{b^2 x^2}{p}.$$

Equations (4.2), (4.3) and Theorem 4.1 ensure that, on an event Ω^x so that $P((\Omega^x)^c) \leq 4e^{-x}$,

$$|U| \leq \kappa' \left(\epsilon \frac{D}{p} + \frac{1}{\epsilon} \left(\frac{v^2 x}{p} + \left(\frac{bx}{\epsilon p} \right)^2 \right) \right),$$

where the constant κ' can be taken equal to $2(\kappa + 4(\ln 2)^{-1})$, for the constant κ defined in Theorem 4.1. Finally, let us recall the following consequence of Bernstein's inequality.

COROLLARY 4.4. *Let $A_0, \dots, A_{p-1}$ be i.i.d random variables valued in a measurable space $(\mathbb{X}, \mathcal{X})$, with common law P . Let μ be a measure on $(\mathbb{X}, \mathcal{X})$ and let $(\psi_\lambda)_{\lambda \in \Lambda}$ be an orthonormal system in $L^2(\mu)$. Let L be a linear functional defined on $L^2(\mu)$ and let $B = \{t = \sum_{\lambda \in \Lambda} a_\lambda L\psi_\lambda, \sum_{\lambda \in \Lambda} a_\lambda^2\}$, $v^2 = \sup_{t \in B} P[(t - Pt)^2]$, $b = \sup_{t \in B} \|t\|_\infty$. Let u be a linear combination of $(\psi_\lambda)_{\lambda \in \Lambda}$ and let $\eta > 0$. For all $x > 0$,*

$$\mathbb{P} \left(\nu_A(Lu) > \frac{\eta}{2} \|u\|^2 + \frac{1}{\eta} \left(\frac{2v^2 x}{p} + \frac{b^2 x^2}{9p^2} \right) \right) \leq e^{-x}.$$

Proof of Corollary 4.4. From Bernstein's inequality, for all $x > 0$, there exists Ω^x satisfying $P(\Omega^x) \geq 1 - e^{-x}$ such that, on Ω^x ,

$$\nu_A(Lu) \leq \sqrt{2 \frac{\text{Var}(Lu(A_1))x}{p}} + \frac{\|Lu\|_\infty x}{3p}.$$

As $L(u/\|u\|)$ belongs to B , we have

$$\|Lu\|_\infty \leq \|u\| b, \quad \text{Var}(Lu(A_1)) \leq \|u\|^2 v^2.$$

Hence, on Ω^x , for all $\eta > 0$, we have

$$\begin{aligned} \nu_A(Lu) &\leq \|u\| \left(\sqrt{2 \frac{v^2 x}{p}} + \frac{bx}{3p} \right) \leq \frac{\eta}{2} \|u\|^2 + \frac{1}{2\eta} \left(\sqrt{2 \frac{v^2 x}{p}} + \frac{bx}{3p} \right)^2 \\ &\leq \frac{\eta}{2} \|u\|^2 + \frac{1}{\eta} \left(2 \frac{v^2 x}{p} + \frac{b^2 x^2}{9p^2} \right). \quad \square \end{aligned}$$

Let us introduce here some notations. For all m in $\mathcal{M}_n$, let $B_m = \{t \in S_m, \|t\|^2 \leq 1\}$,

$$v_{A,m}^2 = q \sup_{t \in B_m} \mathbb{E} \left[(L_q t(A_1) - Pt)^2 \right], \quad b_m = \sup_{t \in B_m} \|t\|_\infty,$$

$$\epsilon_n(p, q) = \sqrt{\ln n} \left(\sup_{m \in \mathcal{M}_n} \left\{ \left(\frac{v_{A,m}^2}{R_{A,m}} \right)^2 \vee \frac{qb_m^2}{pR_{A,m}} \right\} \right)^{1/4}.$$

LEMMA 4.5. *Let $A_0^*, \dots, A_{p-1}^*$ be i.i.d random variables valued in $\mathbb{R}^q$, with $2pq = n$. Let $\mathcal{M}_n$ be a collection of models satisfying Assumptions **H1**, **H2** and let $(p^*(m))_{m \in \mathcal{M}_n}$, $(p_W^*(m))_{m \in \mathcal{M}_n}$, $(\delta^*(m, m'))_{(m, m') \in (\mathcal{M}_n)^2}$, $(D_{A,m})_{m \in \mathcal{M}_n}$, $(R_{A,m})_{m \in \mathcal{M}_n}$ be the associated collections defined in (3.1). Let us assume that $\epsilon_n(p, q)$ is finite. There exist constants κ which may vary from line to line, and an event $\Omega_n^{(2)}$ satisfying $\mathbb{P}(\Omega_n^{(2)}) \geq 1 - \kappa n^{-2}$ such that, on $\Omega_n^{(2)}$, for all m, m' in $\mathcal{M}_n$,*

$$(4.4) \quad \left| p^*(m) - \frac{2D_{A,m}}{n} \right| \leq \kappa \epsilon_n(p, q) \frac{R_{A,m}}{n},$$

$$(4.5) \quad |p^*(m) - p_W^*(m)| \leq \kappa \epsilon_n(p, q) \frac{R_{A,m}}{n},$$

$$(4.6) \quad \delta^*(m, m') \leq \kappa \epsilon_n(p, q) \frac{R_{A,m} \vee R_{A,m'}}{n}.$$

Proof of Lemma 4.5. First, remark that, from Cauchy-Schwarz inequality

$$p^*(m) = \sup_{t \in B_m} (\nu_A(t))^2, \quad \frac{2D_{A,m}}{n} = \mathbb{E}(p^*(m)).$$

We apply Theorem 4.1 to the class $S = \{L_q t, t \in B_m\}$. For all t in S , $P[(t - Pt)^2] \leq v_{A,m}^2/q$ and $\|t\|_\infty \leq b_m$, thus, for all $x > 0$, with probability larger than $1 - 2e^{-x}$, for all $\epsilon > 0$

$$\begin{aligned} \left| p^*(m) - \frac{2D_{A,m}}{n} \right| &\leq \kappa \left(\epsilon \frac{D_{A,m}}{n} + \frac{1}{\epsilon} \left(\frac{v_{A,m}^2 x}{n} + \left(\frac{b_m x}{\epsilon p} \right)^2 \right) \right) \\ &\leq \kappa \frac{R_{A,m}}{n} \left(\epsilon + \frac{1}{\epsilon} \frac{v_{A,m}^2 x}{R_{A,m}} + \frac{1}{\epsilon^3} \frac{q b_m^2 x^2}{p R_{A,m}} \right). \end{aligned}$$

By definition of $\epsilon_n(p, q)$, for all $x > 0$, with probability larger than $1 - 2e^{-x}$, for all $\epsilon > 0$

$$(4.7) \quad \left| p^*(m) - \frac{2D_{A,m}}{n} \right| \leq \kappa \frac{R_{A,m}}{n} \left(\epsilon + \frac{\epsilon_n^2(p, q)}{\epsilon} \frac{x}{\ln n} + \frac{\epsilon_n^4(p, q)}{\epsilon^3} \frac{x^2}{(\ln n)^2} \right).$$

Taking $x = \ln(|\mathcal{M}_n|n^2)$, $\epsilon = \kappa_o \epsilon_n(p, q)$, the constant κ_o being chosen so that, for all $n, \epsilon \leq 1$ and using a union bound yield (4.4).

From (2.1) applied with $(t_\lambda)_{\lambda \in \Lambda} = (L_q(\psi_\lambda))_{\lambda \in \Lambda_m}$, $A_i = A_i^*$,

$$p^*(m) - p_W^*(m) = U_m^*.$$

From Corollary 4.3 applied with $(t_\lambda)_{\lambda \in \Lambda} = (L_q(\psi_\lambda))_{\lambda \in \Lambda_m}$, for all $x > 0$, there exists Ω^x such that $P(\Omega^x) \geq 1 - 4e^{-x}$ and on Ω^x , for all $0 < \epsilon \leq 1$,

$$|U_m^*| \leq \kappa' \left(\epsilon \frac{2D_{A,m}}{n} + \frac{1}{\epsilon} \frac{v_{A,m}^2 x}{n} + \frac{1}{\epsilon^3} \frac{b_m^2 x^2}{p^2} \right).$$

By definition of $\epsilon_n(p, q)$, we deduce that

$$(4.8) \quad |U_m^*| \leq \kappa' \frac{R_{A,m}}{n} \left(\epsilon + \frac{\epsilon_n^2(p, q)}{\epsilon} \frac{x}{\ln n} + \frac{\epsilon_n^4(p, q)}{\epsilon^3} \frac{x^2}{(\ln n)^2} \right).$$

Taking $x = \ln(|\mathcal{M}_n|n^2)$, $\epsilon = \kappa_o \epsilon_n(p, q)$, the constant κ_o being chosen so that, for all $n, \epsilon \leq 1$ and using a union bound yield (4.5).

In order to prove (4.6), we use Corollary 4.4 to the class $S_m + S_{m'}$, with $L = L_q$ and the function $u = s_m - s_{m'}$. It comes from Assumption **H1** that, for all t in $S_m + S_{m'}$ such that $\|t\| \leq 1$,

$$P[(t - Pt)^2] \leq 4 \frac{\kappa_A^2}{q} (v_{A,m}^2 + v_{A,m'}^2), \quad \|t\|_\infty^2 \leq 4 \kappa_A^2 (b_m^2 + b_{m'}^2).$$

Hence, using Corollary 4.4 and the inequality $\|s_m - s_{m'}\|^2 \leq 2(\|s - s_{m'}\|^2 + \|s - s_m\|^2) \leq 2n^{-1}(R_{A,m} + R_{A,m'})$, there exists a constant κ such that, with probability larger than $1 - 2e^{-x}$,

(4.9)

$$\nu_A(s_m - s_{m'}) \leq \left(\frac{1}{\eta} + \kappa \eta \epsilon_n^2(p, q) \right) \left(\frac{R_{A,m}}{n} + \frac{R_{A,m'}}{n} \right) \left(\frac{x}{\ln n} + \frac{x^2}{(\ln n)^2} \right).$$

We apply this inequality with $\eta^{-1} = \epsilon_n(p, q)$ and $x = (2 + 2\alpha_{\mathcal{M}}) \ln n$ and we obtain that, with probability larger than $1 - n^{-2-2\alpha_{\mathcal{M}}}$, for all m, m' in $\mathcal{M}_n$,

$$\nu_A(s_m - s_{m'}) \leq \kappa \epsilon_n^2(p, q) \left(\frac{R_{A,m}}{n} + \frac{R_{A,m'}}{n} \right).$$

A union bound concludes the proof.

LEMMA 4.6. *Let $A_0^*, \dots, A_{p-1}^*$ be i.i.d random variables valued in $\mathbb{R}^q$, with $2pq = n$. Let $\mathcal{M}_n$ be a collection of wavelet spaces $[\mathbf{W}]$ and let $(p^*(m))_{m \in \mathcal{M}_n}$, $(p_W^*(m))_{m \in \mathcal{M}_n}$, $(\delta^*(m, m'))_{(m, m') \in (\mathcal{M}_n)^2}$, $(D_{A,m})_{m \in \mathcal{M}_n}$, $(R_{A,m})_{m \in \mathcal{M}_n}$ be the associated collections defined in (3.1). Let us assume that $\epsilon_n(p, q)$ is finite. There exist constants κ_1, κ_2 which may vary from line to line such that*

$$(4.10) \quad \mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \left(p^*(m) - \frac{2D_{A,m}}{n} - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n} \right)_+ \right) \leq \frac{\kappa_2}{n}.$$

$$(4.11) \quad \mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \left(\frac{2D_{A,m}}{n} - p^*(m) - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n} \right)_+ \right) \leq \frac{\kappa_2}{n}.$$

$$(4.12) \quad \mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \left(p^*(m) - p_W^*(m) - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n} \right)_+ \right) \leq \frac{\kappa_2}{n}.$$

$$(4.13) \quad \mathbb{E} \left(\sup_{m \in \mathcal{M}_n} \left(p_W^*(m) - p^*(m) - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n} \right)_+ \right) \leq \frac{\kappa_2}{n}.$$

$$(4.14) \quad \mathbb{E} \left(\sup_{m, m' \in \mathcal{M}_n} \left(\delta^*(m, m') - \kappa_1 \epsilon_n(p, q) \left(\frac{R_{A,m} \vee R_{A,m'}}{n} \right) \right)_+ \right) \leq \frac{\kappa_2}{n}.$$

Proof: Let us denote by $\mathcal{M}$ the set $\mathcal{M}_n$ or $\mathcal{M}_n^2$ and for all $\overline{m}$ in $\mathcal{M}$, let

$$(4.15) \quad f_1^*(\overline{m}, \kappa_1) = p^*(m) - \frac{2D_{A,m}}{n} - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n},$$

$$(4.16) \quad f_2^*(\overline{m}, \kappa_1) = \frac{2D_{A,m}}{n} - p^*(m) - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n},$$

$$(4.17) \quad f_3^*(\overline{m}, \kappa_1) = p^*(m) - p_W^*(m) - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n},$$

$$(4.18) \quad f_4^*(\overline{m}, \kappa_1) = p_W^*(m) - p^*(m) - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m}}{n},$$

$$(4.19) \quad f_5^*(\overline{m}, \kappa_1) = \delta^*(m, m') - \kappa_1 \epsilon_n(p, q) \frac{R_{A,m} \vee R_{A,m'}}{n}.$$

Fact 8 For all $i = 1, \dots, 5$, we have

$$\mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} (f_i^*(\overline{m}, \kappa_1))_+ \right) \leq \sum_{\overline{m} \in \mathcal{M}} \mathbb{E} ((f_i^*(\overline{m}, \kappa_1))_+) = \sum_{\overline{m} \in \mathcal{M}} \int_0^\infty \mathbb{P}(f_i^*(\overline{m}, \kappa_1) > x) dx.$$

If $\mathcal{M} = \mathcal{M}_n$, for all $\overline{m} = m$ in $\mathcal{M}$, let $R_{A,\overline{m}} = R_{A,m}$. If $\mathcal{M} = \mathcal{M}_n^2$, for all $\overline{m} = (m, m')$ in $\mathcal{M}$, let $R_{A,\overline{m}} = R_{A,m} \vee R_{A,m'}$.

Fact 9 There exist κ_1, κ_2 such that, for all $y > 0$, for all $i = 1, \dots, 5$, for all $\overline{m}$ in $\mathcal{M}$

$$\mathbb{P} \left(f_i^*(\overline{m}, \kappa_1) > \kappa_1 \epsilon_n(p, q) \frac{R_{A,\overline{m}}}{n} (y + y^2) \right) \leq \frac{\kappa_2}{n^2} e^{-y}.$$

Proof: When $i \in \{1, 2\}$, Fact 9 follows from (4.7) applied with $x = y + \ln(|\mathcal{M}_n|n^2)$, $\epsilon = \kappa_o \epsilon_n(p, q)$, the constant κ_o being chosen so that, for all n , $\epsilon \leq 1$. When $i \in \{3, 4\}$, Fact 9 follows from (4.8) applied with $x = y + \ln(|\mathcal{M}_n|n^2)$, $\epsilon = \kappa_o \epsilon_n(p, q)$, with the same constant κ_o . When $i = 5$, Fact 9 follows from (4.9) applied with $x = y + \ln(|\mathcal{M}_n^2|n^2)$, $\epsilon = \kappa_o \epsilon_n(p, q)$, with the same constant κ_o .

Fact 10 Assume that there exist $a_{\overline{m}}, \kappa$ such that

$$\mathbb{P}(f(\overline{m}) > a_{\overline{m}}(x + x^2)) \leq \kappa e^{-x}.$$

There exists a constant κ_1 such that

$$\int_0^\infty \mathbb{P}(f(\overline{m}) > x) dx \leq 3\kappa a_{\overline{m}}.$$

Proof We use the change of variables $x = a_{\overline{m}}(y + y^2)$ in the integral, it gives

$$\int_0^\infty \mathbb{P}(f(\overline{m}) > x) dx \leq \kappa \int_0^\infty e^{-y} a_{\overline{m}} (1 + 2y) dy = 3\kappa a_{\overline{m}}.$$

Facts 8, 9, 10 together give that for all i in $\{1, \dots, 5\}$,

$$\mathbb{E} \left(\sup_{\bar{m} \in \mathcal{M}} (f_i^*(\bar{m}, \kappa_1))_+ \right) \leq \frac{\kappa \epsilon_n(p, q)}{n^2} \sum_{\bar{m} \in \mathcal{M}} \frac{R_{A, \bar{m}}}{n}.$$

It comes from Lemma 5.2 below that, in the collection $[\mathbf{W}]$, for all m in $\mathcal{M}_n$, $R_{A, m} \leq \kappa n$ for some constant κ , hence, $\sum_{\bar{m} \in \mathcal{M}} R_{A, \bar{m}} \leq \kappa n (\ln n)^2 / \ln 2$, which conclude the proof. $\square$

Let us now explain why our Assumptions ensure that, in both the β -mixing and the τ -mixing cases, $\epsilon_n(p, q) \leq \kappa (\ln n)^{-1/2}$.

5. Computation of $\epsilon_n(p, q)$.

5.1. *The β -mixing case.* We recall the following inequality for β -mixing processes (see for example Comte and Merlevède (2002) inequalities (6.2) and (6.3)). Let $(X_n)_{n \in \mathbb{Z}}$ be β -mixing data. There exists a function ν satisfying, for all p, q in $\mathbb{N}$,

$$(5.1) \quad \nu = \sum_{l \geq 1} \nu_l, \text{ with } P\nu_q \leq \beta_q \text{ and } P(\nu^p) \leq p \sum_{l \geq 1} (l+1)^{p-1} \beta_l,$$

such that, for all t in $L^2(\mu)$,

$$(5.2) \quad \left| \text{Var} \left(\sum_{i=1}^q t(X_i) \right) - \sum_{i=1}^q \text{Var}(t(X_i)) \right| \leq 4qP(\nu t^2).$$

LEMMA 5.1. *Let $\theta > 1$ and let $(X_n)_{n \in \mathbb{Z}}$ be an arithmetically $[\mathbf{AR}(\theta)]$ β -mixing process satisfying $\mathbf{S}(\beta)$. Let S_m be a linear space satisfying $\mathbf{H3}$, $\mathbf{H4}$. Let p, q such that $2pq = n$, $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$. We have*

$$\frac{c_D}{4} b_m^2 - \|s_m\|^2 \leq D_{A, m} \leq \frac{5}{2} b_m^2 - \|s_m\|^2.$$

$$v_{A, m}^2 \leq \kappa \frac{R_{A, m}}{(\ln n)^2}, \quad \frac{q b_m^2}{p} \leq \kappa \frac{R_{A, m}}{(\ln n)^4}.$$

In particular, $\epsilon_n(p, q) \leq \kappa (\ln n)^{-1/2}$.

Proof: We have, from inequality (5.2),

$$\begin{aligned} & \left| D_{A, m} - \sum_{\lambda \in \Lambda_m} \text{Var}(\psi_\lambda(X_1)) \right| \\ &= \left| q \sum_{\lambda \in \Lambda_m} \text{Var} \left(\frac{1}{q} \sum_{i=1}^q \psi_\lambda(X_i) \right) - \sum_{\lambda \in \Lambda_m} \text{Var}(\psi_\lambda(X_1)) \right| \leq 4P \left(\nu \sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right). \end{aligned}$$

Moreover

$$\sum_{\lambda \in \Lambda_m} \text{Var}(\psi_\lambda(X_1)) = P \left(\sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right) - \sum_{\lambda \in \Lambda_m} (P\psi_\lambda)^2 = P \left(\sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right) - \|s_m\|^2.$$

It comes from Cauchy-Schwarz inequality that $\sum_{\lambda \in \Lambda_m} \psi_\lambda^2 = (\sup_{t \in B_m} t)^2$, hence, from **H4**,

$$c_D b_m^2 \leq P \left(\sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right) \leq b_m^2.$$

Finally,

$$P \left(\nu \sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right) \leq b_m \sqrt{P\nu^2 P \left(\sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right)} \leq \frac{1}{8} P \left(\sum_{\lambda \in \Lambda_m} \psi_\lambda^2 \right) + 2b_m^2 P(\nu^2).$$

We deduce that

$$\left(\frac{c_D}{2} - 8P(\nu^2) \right) b_m^2 - \|s_m\|^2 \leq D_{A,m} \leq \left(\frac{3}{2} + 8P(\nu^2) \right) b_m^2 - \|s_m\|^2.$$

From (5.1) and **S**(β), we have $8P(\nu^2) = 16 \sum_{l \geq 1} (1+l)^1 \beta_l < c_D/4$. As $R_{A,m} \geq \kappa_R (\ln n)^8$

$$b_m^2 \leq \frac{4}{c_D} (D_{A,m} + \|s_m\|^2) \leq \frac{4}{c_D} \left(1 + \frac{\|s_m\|^2}{\kappa_R} \right) R_{A,m}.$$

Let us then denote by $\kappa_\infty = \frac{4}{c_D} \left(1 + \frac{\|s\|^2}{\kappa_R} \right)$, we have obtained that $b_m^2 \leq \kappa_\infty R_{A,m}$. Our definition of p and q yields then

$$\frac{qb_m^2}{p} \leq \kappa \frac{R_{A,m}}{(\ln n)^4}.$$

We also have

$$v_{A,m}^2 = q \sup_{t \in B_m} P[(L_q t - pt)^2].$$

Let t in B_m , we have, from inequality (5.2) and **S**(β)

$$\begin{aligned} \text{Var} \left(\frac{1}{q} \sum_{i=1}^q t(X_i) \right) &\leq \frac{1}{q} P((1+4\nu)t^2) \leq \frac{1}{q} \|t\|_\infty (P|t| + 4\sqrt{P\nu^2 P t^2}) \\ &\leq \frac{2b_m}{q} \sqrt{P t^2} \leq \frac{2\sqrt{\|s\|} b_m^{3/2}}{q} \leq 2\sqrt{\|s\|} \kappa_\infty^{3/4} \frac{R_{A,m}}{R_n^{1/4} q}. \end{aligned}$$

Since $R_n \geq \kappa_R(\ln n)^8$, we deduce that

$$\frac{v_{A,m}^2}{R_{A,m}} \leq \frac{2\sqrt{\|s\|}\kappa_\infty^{3/4}}{\kappa_R^{1/4}}(\ln n)^{-2}.$$

5.2. The τ -mixing case.

LEMMA 5.2. *Let c_τ be the constant defined in Lemma 5.3 and let $C(\phi, \psi) = c_D \kappa_\infty^2 / (8c_\tau A K_\infty K_{BV})$. Let $\theta > 1$ and let $(X_n)_{n \in \mathbb{Z}}$ be an arithmetically $[\mathbf{AR}(\theta)]$ τ -mixing process satisfying $\mathbf{S}(\tau, \mathbf{W})$. Let S_m be a linear space in the collection $[\mathbf{W}]$ satisfying Assumptions **H3**, **H4**. Let p, q such that $2pq = n$, $\sqrt{n}(\ln n)^2/2 \leq p \leq \sqrt{n}(\ln n)^2$. We have*

$$(5.3) \quad \frac{c_D \kappa_\infty^2}{2} 2^{J_m} - \|s_m\|^2 \leq D_{A,m} \leq \left(AK_\infty^2 + \frac{c_D \kappa_\infty^2}{2} \right) 2^{J_m} - \|s_m\|^2.$$

$$v_{A,m}^2 \leq \kappa \frac{R_{A,m}}{(\ln n)^2}, \quad \frac{qb_m^2}{p} \leq \kappa \frac{R_{A,m}}{(\ln n)^4}.$$

In particular, $\epsilon_n(p, q) \leq \kappa(\ln n)^{-1/2}$.

Proof:

Control of b_m^2 : We use the following elementary inequalities:

$$\begin{aligned} \left\| \sum_{(j,k) \in \Lambda_m} \psi_{j,k}^2 \right\|_\infty &\leq AK_\infty^2 2^{J_m} \\ \left\| \sum_{(j,k) \in \Lambda_m} \psi_{j,k}^2 \right\|_\infty &\geq \max_{(j,k) \in \Lambda_m} \|\psi_{j,k}^2\|_\infty \geq \kappa_\infty^2 2^{J_m}. \end{aligned}$$

We obtain that

$$(5.4) \quad \kappa_\infty^2 2^{J_m} \leq b_m^2 \leq AK_\infty^2 2^{J_m}.$$

Control of $v_{A,m}^2$: Let t be a function in B_m . First, we use a simple bound

$$q \text{Var} \left(\frac{1}{q} \sum_{i=1}^q t(X_i) \right) \leq 2 \sum_{l=1}^q |\text{Cov}(t(X_1), t(X_l))|.$$

We use the coupling lemma obtained in [Dedecker and Prieur \(2005\)](#) Section 7.1, to define random variables X_l^* independent of X_1 , such that

$$\mathbb{E}(|X_l - X_l^*|) \leq \tau_{l-1}.$$

$$\begin{aligned}
|\text{Cov}(t(X_1), t(X_l))| &= |\text{Cov}(t(X_1), t(X_l) - t(X_l^*))| \\
&\leq \sqrt{\text{Var}(t(X_1)) \mathbb{E}[(t(X_l) - t(X_l^*))^2]} \\
&\leq \sqrt{\text{Var}(t(X_1)) 2b_m \mathbb{E}[|t(X_l) - t(X_l^*)|]} \leq \sqrt{\text{Var}(t(X_1)) 2b_m \text{Lip}(t) \tau_{l-1}}
\end{aligned}$$

Moreover, let $a_{j,k} = \int_{\mathbb{R}} t \psi_{j,k} d\mu$, then

$$\begin{aligned}
\text{Lip}(t) &= \sup_{x \neq y \in \mathbb{R}} \frac{|t(x) - t(y)|}{|x - y|} \leq \sum_{j=0}^{J_m} \sup_{x \neq y \in \mathbb{R}} \sum_{k \in \mathbb{Z}} |a_{j,k}| \frac{|\psi_{j,k}(x) - \psi_{j,k}(y)|}{|x - y|} \\
(5.5) \quad &\leq 2AK_L \sum_{j=0}^{J_m} 2^{3j/2} \sup_{k \in \mathbb{Z}} |a_{j,k}|.
\end{aligned}$$

The last inequality holds since, for all x, y in $\mathbb{R}$ there is less than $2A$ indices k in $\mathbb{Z}$ such that $|\psi_{j,k}(x) - \psi_{j,k}(y)| \neq 0$. Since t belongs to B_m , $\sum_{(j,k) \in \Lambda_m} a_{j,k}^2 \leq 1$, in particular, for all j , $\sup_{k \in \mathbb{Z}} |a_{j,k}| \leq 1$. Thus, there exists a constant c such that $\text{Lip}(t) \leq c2^{3J_m/2}$. Hence, there exists a constant c such that, for all t in B_m and all l in $\mathbb{N}^*$

$$|\text{Cov}(t(X_1), t(X_l))| \leq c2^{5J_m/4} \sqrt{\tau_{l-1}}.$$

Remark that we also have

$$|\text{Cov}(t(X_1), t(X_l))| \leq \|t\|_{\infty} \|t\| \|s\| \leq c2^{J_m/2}.$$

Let $u = 3/(1 + \theta)$, there exist constants c , which may vary from line to line such that

$$\begin{aligned}
\sum_{l=1}^q |\text{Cov}(t(X_1), t(X_l))| &\leq c2^{J_m/2} \sum_{l=1}^{\infty} (2^{3J_m/4} \sqrt{\tau_{l-1}} \wedge 1) \\
&\leq c2^{J_m/2} \sum_{l=1}^{\infty} (2^{3J_m/4} l^{-(1+\theta)/2} \wedge 1) \\
&\leq c2^{J_m/2} \left(\sum_{l=1}^{2^{uJ_m/2}} 1 + \sum_{l=2^{uJ_m/2}}^{\infty} 2^{3J_m/4} l^{-(1+\theta)/2} \right) \leq c2^{\frac{J_m}{2}(1+u)}.
\end{aligned}$$

Since $\theta > 5$, $u < 1/2$ and we have obtained that

$$(5.6) \quad v_{A,m}^2 \leq \kappa 2^{3J_m/4}.$$

Control of $D_{A,m}$: Let us recall the following lemma, obtained in [Lerasle \(2009a\)](#) as a consequence of the covariance inequality for τ -mixing sequences proved by [Dedecker and Prieur \(2005\)](#).

LEMMA 5.3. *Let X, Y be two identically distributed real valued random variables, with common density s in $L^2(\mu)$. There exist a constant c_τ and a random variable $\nu(\sigma(X), Y)$ satisfying*

$$\mathbb{E}(\nu(\sigma(X), Y)) = c_\tau \|s\|^{2/3} (\tau(\sigma(X), Y))^{1/3}$$

such that, for all Lipschitz functions g and all h in BV

$$(5.7) \quad \begin{aligned} |\text{Cov}(g(X), h(Y))| &\leq \|h\|_{BV} \mathbb{E}(|g(X)|\nu(\sigma(X), Y)) \\ &\leq c_\tau \|h\|_{BV} \|g\|_\infty \|s\|^{2/3} (\tau(\sigma(X), Y))^{1/3}. \end{aligned}$$

It comes from this Lemma and inequalities (3.5, 3.6, 3.7) that

$$\begin{aligned} &|D_{A,m} - \sum_{(j,k) \in \Lambda_m} \text{Var}(\psi_{j,k}(X_1))| \\ &\leq 2 \sum_{(j,k) \in \Lambda_m} \sum_{l=2}^q (q+1-l) |\text{Cov}(\psi_{j,k}(X_1), \psi_{j,k}(X_l))| \\ &\leq \frac{2}{q} \sum_{j=0}^{J_m} \sum_{k \in \mathbb{Z}} \sum_{l=2}^q \|\psi_{j,k}\|_{BV} \mathbb{E}(|\psi_{j,k}(X_1)|\nu(\sigma(X_1), X_l)) \\ &\leq 2c_\tau K_{BV} \|s\|^{2/3} \sum_{j=0}^{J_m} 2^{j/2} \left\| \sum_{k \in \mathbb{Z}} |\psi_{j,k}| \right\|_\infty \sum_{l=2}^q \tau_{l-1}^{1/3} \\ &\leq 4 \left(c_\tau A K_\infty K_{BV} \|s\|^{2/3} \sum_{l=1}^\infty \tau_l^{1/3} \right) 2^{J_m} \leq \frac{c_D \kappa_\infty}{2} 2^{J_m}. \end{aligned}$$

The last inequality comes from $\mathbf{S}(\tau, \mathbf{W})$. Since $\sum_{(j,k) \in \Lambda_m} (P\psi_{j,k}(X_1))^2 = \|s_m\|^2$, we deduce that

$$-\frac{c_D \kappa_\infty^2}{2} 2^{J_m} \leq D_{A,m} - P \left(\sum_{(j,k) \in \Lambda_m} \psi_{j,k}^2(X_1) \right) + \|s_m\|^2 \leq \frac{c_D \kappa_\infty^2}{2} 2^{J_m}$$

We always have $P \left(\sum_{(j,k) \in \Lambda_m} \psi_{j,k}^2(X_1) \right) = P \left(\sup_{t \in B_m} t^2 \right) \leq b_m^2 \leq A K_\infty^2 2^{J_m}$. From **H4**,

$$P \left(\sum_{(j,k) \in \Lambda_m} \psi_{j,k}^2(X_1) \right) = P \left(\sup_{t \in B_m} t^2 \right) \geq c_D b_m^2 \geq c_D \kappa_\infty^2 2^{J_m}.$$

We obtain finally

$$(5.8) \quad \frac{c_D \kappa_\infty^2}{2} 2^{J_m} - \|s_m\|^2 \leq D_{A,m} \leq \left(AK_\infty^2 + \frac{c_D \kappa_\infty^2}{2} \right) 2^{J_m} - \|s_m\|^2.$$

Conclusion of the proof: It comes from inequalities (5.6) and (5.8) that

$$v_{A,m}^2 \leq \kappa(D_{A,m} + \|s\|^2)^{3/4} \leq \kappa(R_{A,m} + \|s\|^2)^{3/4}.$$

As $R_{A,m} \geq \kappa_R(\ln n)^8$, this implies that

$$v_{A,m}^2 \leq \kappa \frac{R_{A,m}}{R_n^{1/4}} \leq \kappa \frac{R_{A,m}}{(\ln n)^2}.$$

Now, from inequalities (5.4) and (5.8), we have

$$b_m^2 \leq \kappa(D_{A,m} + \|s\|^2) \leq \kappa' R_{A,m}.$$

Hence, our choice of p and q ensures that

$$\frac{qb_m^2}{p} \leq \kappa \frac{R_{A,m}}{(\ln n)^2}.$$

6. Conclusion of the proofs.

6.1. *Proof of Lemma 1.1.* Let us use the notations $f_i^*(\overline{m}, \kappa_1)$ defined in (4.15-4.19). For all $i = 1, \dots, 5$,

$$\begin{aligned} & \mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} (f_i(\overline{m}, \kappa_1))_+ \right) \\ & \leq \mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} |f_i(\overline{m}, \kappa_1) - f_i^*(\overline{m}, \kappa_1)| \right) + \mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} (f_i^*(\overline{m}, \kappa_1))_+ \right). \end{aligned}$$

It comes from Lemma 3.4 that, for all $i = 1, \dots, 5$,

$$\mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} |f_i(\overline{m}, \kappa_1) - f_i^*(\overline{m}, \kappa_1)| \right) \leq (4 \vee 8/p) \tau_q M C_n.$$

From Lemma 3.6 that $\tau_q M C_n \leq \kappa n^{-1}$ since $\theta > 5$. Moreover, Lemmas 4.6 and 5.2 ensure that $\mathbb{E} \left(\sup_{\overline{m} \in \mathcal{M}} (f_i^*(\overline{m}, \kappa_1))_+ \right) \leq \frac{\kappa}{n}$.

6.2. *Proof of Lemma 1.2.* For all $i = 1, \dots, 5$ and all $\overline{m}$ in $\mathcal{M}$, let $f_i^*(\overline{m})$, $f_i(\overline{m})$ be the random variables defined in (4.15-4.19) and (1.1-1.5) for $\kappa_1 = 0$. Let Ω^* be the set defined by Lemma 3.3. From Lemma 3.5, we have $P((\Omega^*)^c) \leq \kappa_2(\ln n)^{2(1+\theta)}n^{-\theta/2}$ and, on Ω^* ,

$$\forall i = 1, \dots, 5, \forall \overline{m} \in \mathcal{M}, |f_i(\overline{m})| = |f_i^*(\overline{m})|.$$

From Lemmas 4.5 and 5.1, there exists a set Ω_* , with $P(\Omega_*^c) \leq \kappa n^{-2}$ such that, on Ω_* ,

$$\forall i = 1, \dots, 5, \forall \overline{m} \in \mathcal{M}, |f_i^*(\overline{m})| \leq \kappa_1 \varepsilon_n \frac{R_{A, \overline{m}}}{n}.$$

Hence, on $\Omega_* \cap \Omega^*$, the conclusion of the lemma holds.

References.

- BOUSQUET, O. (2002). A Bennett concentration inequality and its application to suprema of empirical processes. *C. R. Math. Acad. Sci. Paris* **334** 495–500. [MR1890640 \(2003f:60039\)](#)
- COMTE, F. and MERLEVÈDE, F. (2002). Adaptive estimation of the stationary density of discrete and continuous time mixing processes. *ESAIM Probab. Statist.* **6** 211–238 (electronic). New directions in time series analysis (Luminy, 2001). [MR1943148 \(2004b:62105\)](#)
- DEDECKER, J. and PRIEUR, C. (2005). New dependence coefficients. Examples and applications to statistics. *Probab. Theory Related Fields* **132** 203–236. [MR2199291 \(2007b:62081\)](#)
- KLEIN, T. and RIO, E. (2005). Concentration around the mean for maxima of empirical processes. *Ann. Probab.* **33** 1060–1077. [MR2135312 \(2006c:60022\)](#)
- LERASLE, M. (2009a). Adaptive density estimation of stationary β -mixing and τ -mixing sequences. *Mathematical Methods of statistics*.
- LERASLE, M. (2009b). Optimal model selection in density estimation. *arXiv:0910.1654*.
- VIENNET, G. (1997). Inequalities for absolutely regular sequences: application to density estimation. *Probab. Theory Related Fields* **107** 467–492. [MR1440142 \(98f:62113\)](#)

E-MAIL: lerasle@gmail.com